

## ARG42313 anti-CD279 / PD-1 antibody [EH12.2H7] (FITC)

Package: 50 tests

Store at: 4°C

### Summary

Product Description	FITC-conjugated Mouse Monoclonal antibody [EH12.2H7] recognizes CD279 / PD-1
Tested Reactivity	Hu, NHuPrm
Tested Application	FACS
Specificity	The mouse monoclonal antibody EH12.2H7 recognizes an extracellular epitope of CD279 / PD-1 (programmed cell death 1), a 55 kDa type I transmembrane protein expressed above all during T cell development, on activated T cells, activated B cells, and activated monocytes.
Host	Mouse
Clonality	Monoclonal
Clone	EH12.2H7
Isotype	IgG1, kappa
Target Name	CD279 / PD-1
Species	Human
Immunogen	Human CD279.
Conjugation	FITC
Alternate Names	hPD-1; CD279; PD-1; Protein PD-1; CD antigen CD279; PD1; hSLE1; SLEB2; Programmed cell death protein 1; hPD-1

### Application Instructions

Application table	Application	Dilution
	FACS	4 µl / 100 µl of whole blood or 10 <sup>6</sup> cells

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


### Properties

Form	Liquid
Purification	Purified
Buffer	PBS and 15 mM Sodium azide.
Preservative	15 mM Sodium azide
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	PDCD1
Gene Full Name	programmed cell death 1
Background	This gene encodes a cell surface membrane protein of the immunoglobulin superfamily. This protein is expressed in pro-B-cells and is thought to play a role in their differentiation. In mice, expression of this gene is induced in the thymus when anti-CD3 antibodies are injected and large numbers of thymocytes undergo apoptosis. Mice deficient for this gene bred on a BALB/c background developed dilated cardiomyopathy and died from congestive heart failure. These studies suggest that this gene product may also be important in T cell function and contribute to the prevention of autoimmune diseases. [provided by RefSeq, Jul 2008]
Function	<p>Inhibitory receptor on antigen activated T-cells that plays a critical role in induction and maintenance of immune tolerance to self (PubMed:21276005). Delivers inhibitory signals upon binding to ligands CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed:21276005). Following T-cell receptor (TCR) engagement, PDCD1 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the ITSM motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKCtheta and CD247/CD3zeta (By similarity).</p> <p>The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed:28951311). The interaction with CD274/PDCD1L1 inhibits cytotoxic T lymphocytes (CTLs) effector function (PubMed:28951311). The blockage of the PDCD1-mediated pathway results in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (PubMed:22658127, PubMed:25034862, PubMed:25399552). [UniProt]</p>
Highlight	<p>Related products: <a href="#">PD-1 antibodies</a>; <a href="#">PD-1 ELISA Kits</a>; <a href="#">PD-1 Duos / Panels</a>; <a href="#">Anti-Mouse IgG secondary antibodies</a>;</p> <p>Related news: <a href="#">Examining CTL/NK-mediated cytotoxicity by ELISA</a></p>
Calculated Mw	32 kDa
Cellular Localization	Membrane; Single-pass type I membrane protein. [UniProt]

## Images



ARG42313 anti-CD279 / PD-1 antibody [EH12.2H7] (FITC) FACS image

Flow Cytometry: Human PHA stimulated peripheral blood mononuclear cells stained with ARG42313 anti-CD279 / PD-1 antibody [EH12.2H7] (FITC) at 4  $\mu$ l /  $10^6$  cells in 100  $\mu$ l of cell suspension.



ARG42313 anti-CD279 / PD-1 antibody [EH12.2H7] (FITC) FACS image

Flow Cytometry: Separation of Human CD279 positive cells (red-filled) from cellular debris (black-dashed). Human PHA stimulated peripheral blood mononuclear cells stained with ARG42313 anti-CD279 / PD-1 antibody [EH12.2H7] (FITC) at 4  $\mu$ l / 10<sup>6</sup> cells in 100  $\mu$ l of cell suspension.