

## ARG42480 anti-COPG2 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes COPG2
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	COPG2
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 542-871 of Human COPG2 (NP_036265.3).
Conjugation	Un-conjugated
Alternate Names	Gamma-2-COP; Gamma-2-coat protein; gamma-2-COP; Coatomer subunit gamma-2; 2-COP

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	OVCAR3	
Observed Size	~ 100 kDa	


### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation


Gene Symbol	COPG2
Gene Full Name	coatomer protein complex, subunit gamma 2
Function	The coatomer is a cytosolic protein complex that binds to dilysine motifs and reversibly associates with Golgi non-clathrin-coated vesicles, which further mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins; the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors (By similarity). [UniProt]
Calculated Mw	98 kDa
Cellular Localization	Cytoplasm, cytosol. Golgi apparatus membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, COPI-coated vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Note=The coatomer is cytoplasmic or polymerized on the cytoplasmic side of the Golgi, as well as on the vesicles/buds originating from it. Tends to be more abundant in the trans-Golgi network compared to the cis-Golgi. [UniProt]

## Images



ARG42480 anti-COPG2 antibody ICC/IF image

Immunofluorescence: U2OS cells stained with ARG42480 anti-COPG2 antibody at 1:100 dilution.



ARG42480 anti-COPG2 antibody WB image

Western blot: 25 µg of OVCAR3 cell lysate stained with ARG42480 anti-COPG2 antibody at 1:1000 dilution.