

**ARG42645**  
anti-GAD67 antibodyPackage: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes GAD67
Tested Reactivity	Hu, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	GAD67
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 529-543 of Human GAD67. (PQRREKLHKVAPKIK)
Conjugation	Un-conjugated
Alternate Names	EC 4.1.1.15; Glutamate decarboxylase 67 kDa isoform; Glutamate decarboxylase 1; 67 kDa glutamic acid decarboxylase; GAD; SCP; GAD-67; CPSQ1

### Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 67 kDa	


### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.9% NaCl, 0.05% Sodium azide, 0.05% Thimerosal and 5% BSA.
Preservative	0.05% Sodium azide and 0.05% Thimerosal
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Gene Symbol	GAD1
Gene Full Name	glutamate decarboxylase 1 (brain, 67kDa)
Background	This gene encodes one of several forms of glutamic acid decarboxylase, identified as a major autoantigen in insulin-dependent diabetes. The enzyme encoded is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantigen and an autoreactive T cell target in insulin-dependent diabetes. This gene may also play a role in the stiff man syndrome. Deficiency in this enzyme has been shown to lead to pyridoxine dependency with seizures. Alternative splicing of this gene results in two products, the predominant 67-kD form and a less-frequent 25-kD form. [provided by RefSeq, Jul 2008]
Function	Catalyzes the production of GABA. [UniProt]
Calculated Mw	67 kDa

## Images



ARG42645 anti-GAD67 antibody WB image

Western blot: 50 µg of samples under reducing conditions. Rat brain, Rat testis, MCF7, MDA-MB-231, HeLa, SMMC-7721 and COLO320 whole cell lysates stained with ARG42645 anti-GAD67 antibody at 0.5 µg/ml dilution, overnight at 4°C.