

ARG42946 anti-NLRP2 / NALP2 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes NLRP2 / NALP2
Tested Reactivity	Hu, Ms
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	NLRP2 / NALP2
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 1-220 of Human NLRP2 / NALP2 (NP_060322.1).
Conjugation	Un-conjugated
Alternate Names	PAN1; PYPAF2; PYRIN-containing APAF1-like protein 2; NALP2; NACHT, LRR and PYD domains-containing protein 2; CLR19.9; NBS1; Nucleotide-binding site protein 1; PYRIN domain and NACHT domain-containing protein 1

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A549	
Observed Size	~ 120 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation


Gene Symbol	NLRP2
Gene Full Name	NLR family, pyrin domain containing 2
Background	This gene is a member of the nucleotide-binding and leucine-rich repeat receptor (NLR) family, and is predicted to contain an N-terminal pyrin effector domain (PYD), a centrally-located nucleotide-binding and oligomerization domain (NACHT) and C-terminal leucine-rich repeats (LRR). Members of this gene family are thought to be important regulators of immune responses. This gene product interacts with components of the I κ B kinase (IKK) complex, and can regulate both caspase-1 and NF- κ B (nuclear factor kappa-light-chain-enhancer of activated B cells) activity. The pyrin domain is necessary and sufficient for suppression of NF- κ B activity. An allelic variant (rs147585490) has been found that is incapable of blocking the transcriptional activity of NF- κ B. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2016]
Function	Suppresses TNF- and CD40-induced NFKB1 activity at the level of the IKK complex, by inhibiting NFKBIA degradation induced by TNF. When associated with PYCARD, activates CASP1, leading to the secretion of mature proinflammatory cytokine IL1B. May be a component of the inflammasome, a protein complex which also includes PYCARD, CARD8 and CASP1 and whose function would be the activation of proinflammatory caspases. [UniProt]
Calculated Mw	121 kDa
Cellular Localization	Cytoplasm. [UniProt]

Images



ARG42946 anti-NLRP2 / NALP2 antibody ICC/IF image

Immunofluorescence: U2OS cells stained with ARG42946 anti-NLRP2 / NALP2 antibody at 1:100 dilution.



A549

ARG42946 anti-NLRP2 / NALP2 antibody WB image

Western blot: 25 μ g of A549 cell lysate stained with ARG42946 anti-NLRP2 / NALP2 antibody at 1:1000 dilution.