

ARG42947 anti-PTPN1 / PTP1B antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes PTPN1 / PTP1B
Tested Reactivity	Hu, Ms, Rat
Tested Application	FACS, IHC-Fr, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PTPN1 / PTP1B
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 4-30 of Human PTPN1 / PTP1B. (EKEFEQIDKSGSWAAIYQDIRHEASDF)
Conjugation	Un-conjugated
Alternate Names	PTP1B; EC 3.1.3.48; Tyrosine-protein phosphatase non-receptor type 1; PTP-1B; Protein-tyrosine phosphatase 1B

Application Instructions

Application table	Application	Dilution
	FACS	1:150 - 1:500
	IHC-Fr	1:200 - 1:1000
	IHC-P	1:200 - 1:1000
	WB	1:500 - 1:2000
Application Note	IHC-P: Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	50 - 55 kDa	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na ₂ HPO ₄ , 0.9% NaCl, 0.05% Sodium azide and 4% Trehalose.
Preservative	0.05% Sodium azide
Stabilizer	4% Trehalose
Concentration	0.5 mg/ml

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	PTPN1
Gene Full Name	protein tyrosine phosphatase, non-receptor type 1
Background	The protein encoded by this gene is the founding member of the protein tyrosine phosphatase (PTP) family, which was isolated and identified based on its enzymatic activity and amino acid sequence. PTPs catalyze the hydrolysis of the phosphate monoesters specifically on tyrosine residues. Members of the PTP family share a highly conserved catalytic motif, which is essential for the catalytic activity. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP has been shown to act as a negative regulator of insulin signaling by dephosphorylating the phosphotyrosine residues of insulin receptor kinase. This PTP was also reported to dephosphorylate epidermal growth factor receptor kinase, as well as JAK2 and TYK2 kinases, which implicated the role of this PTP in cell growth control, and cell response to interferon stimulation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2013]
Function	Tyrosine-protein phosphatase which acts as a regulator of endoplasmic reticulum unfolded protein response. Mediates dephosphorylation of EIF2AK3/PERK; inactivating the protein kinase activity of EIF2AK3/PERK. May play an important role in CKII- and p60c-src-induced signal transduction cascades. May regulate the EFNA5-EPHA3 signaling pathway which modulates cell reorganization and cell-cell repulsion. May also regulate the hepatocyte growth factor receptor signaling pathway through dephosphorylation of MET. [UniProt]
Calculated Mw	50 kDa
PTM	<p>Oxidized on Cys-215; the Cys-SOH formed in response to redox signaling reacts with the alpha-amido of the following residue to form a sulfenamide cross-link, triggering a conformational change that inhibits substrate binding and activity. The active site can be restored by reduction.</p> <p>Ser-50 is the major site of phosphorylation as compared to Ser-242 and Ser-243. Activated by phosphorylation at Ser-50.</p> <p>S-nitrosylation of Cys-215 inactivates the enzyme activity.</p> <p>Sulfhydration at Cys-215 following endoplasmic reticulum stress inactivates the enzyme activity, promoting EIF2AK3/PERK activity. [UniProt]</p>
Cellular Localization	Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Note=Interacts with EPHA3 at the cell membrane. [UniProt]




ARG42947 anti-PTPN1 / PTP1B antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human placenta tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 µg/ml dilution, overnight at 4°C.




ARG42947 anti-PTPN1 / PTP1B antibody WB image

Western blot: 50 µg of sample under reducing conditions. Rat kidney and Mouse kidney lysates stained with ARG42947 anti-PTPN1 / PTP1B antibody at 0.5 µg/ml dilution, overnight at 4°C.




ARG42947 anti-PTPN1 / PTP1B antibody FACS image

Flow Cytometry: U2OS cells were blocked with 10% normal goat serum and then stained with ARG42947 anti-PTPN1 / PTP1B antibody (blue) at 1 µg/10⁶ cells for 30 min at 20°C, followed by incubation with DyLight[®]488 labelled secondary antibody. Isotype control antibody (green) was Rabbit IgG (1 µg/10⁶ cells) used under the same conditions. Unlabelled sample (red) was also used as a control.




ARG42947 anti-PTPN1 / PTP1B antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human mammary cancer tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 µg/ml dilution, overnight at 4°C.




ARG42947 anti-PTPN1 / PTP1B antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human tonsil tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 μ g/ml dilution, overnight at 4°C.




ARG42947 anti-PTPN1 / PTP1B antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse spleen tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 μ g/ml dilution, overnight at 4°C.




ARG42947 anti-PTPN1 / PTP1B antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Mouse testis tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 μ g/ml dilution, overnight at 4°C.




ARG42947 anti-PTPN1 / PTP1B antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Rat spleen tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 μ g/ml dilution, overnight at 4°C.




ARG42947 anti-PTPN1 / PTP1B antibody IHC-Fr image

Immunohistochemistry: Frozen section of Human placenta tissue. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 µg/ml dilution, overnight at 4°C.



ARG42947 anti-PTPN1 / PTP1B antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human tonsil tissue. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0) for 20 min. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG42947 anti-PTPN1 / PTP1B antibody at 1 µg/ml dilution, overnight at 4°C.



ARG42947 anti-PTPN1 / PTP1B antibody WB image

Western blot: 50 µg of sample under reducing conditions. Human placenta, MCF7, Caco-2, K562, U2OS, PC-3, HL-60 and A549 whole cell lysates stained with ARG42947 anti-PTPN1 / PTP1B antibody at 0.5 µg/ml dilution, overnight at 4°C.