

# Product datasheet

info@arigobio.com

ARG43189 anti-BCL9L antibody

Package: 50 μg Store at: -20°C

## **Summary**

Product Description Rabbit Polyclonal antibody recognizes BCL9L

Tested Reactivity Hu

Tested Application ICC/IF, WB
Host Rabbit
Clonality Polyclonal

Isotype IgG

Target Name BCL9L

Species Human

Immunogen A 20-amino acid peptide within aa. 20-70 of Human BCL9L.

Conjugation Un-conjugated

Alternate Names Protein BCL9-2; BCL9-2; B-cell CLL/lymphoma 9-like protein; DLNB11; BCL9-like protein; B-cell

lymphoma 9-like protein

## **Application Instructions**

Application table	Application	Dilution
	ICC/IF	10 μg/ml
	WB	1 μg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	

### **Properties**

Form Liquid

Purification Affinity purification with immunogen.

Buffer PBS and 0.02% Sodium azide.

Preservative 0.02% Sodium azide

Concentration 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Gene Symbol BCL9L

Gene Full Name B-cell CLL/lymphoma 9-like

Function Transcriptional regulator that acts as an activator. Promotes beta-catenin transcriptional activity. Plays

a role in tumorigenesis. Enhances the neoplastic transforming activity of CTNNB1 (By similarity).

[UniProt]

Calculated Mw 157 kDa


Cellular Localization Nucleus. [UniProt]

# **Images**



#### ARG43189 anti-BCL9L antibody ICC image

Immunocytochemistry: HeLa cells stained with ARG43189 anti-BCL9L antibody at 10  $\mu g/ml$  dilution.



#### ARG43189 anti-BCL9L antibody WB image

Western blot: HeLa cell lysate stained with ARG43189 anti-BCL9L antibody at 1  $\mu$ g/ml dilution, in the absence (left) and the presence of blocking peptide (right).