

# Product datasheet

info@arigobio.com

ARG43545 anti-EGR2 antibody

Package: 100 μl Store at: -20°C

#### **Summary**

Product Description Rabbit Polyclonal antibody recognizes EGR2.

Tested Reactivity Hu, Ms, Rat

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name EGR2

Species Human

Immunogen Synthetic peptide derived from human EGR2

Conjugation Un-conjugated

Alternate Names AT591; CMT1D; CMT4E; KROX20

## **Application Instructions**

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

## **Properties**

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

|--|

www.arigobio.com arigo.nuts about antibodies 1/2

Gene Full Name early growth response 2

Background The protein encoded by this gene is a transcription factor with three tandem C2H2-type zinc fingers.

Defects in this gene are associated with Charcot-Marie-Tooth disease type 1D (CMT1D), Charcot-Marie-Tooth disease type 4E (CMT4E), and with Dejerine-Sottas syndrome (DSS). Multiple transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Oct 2008]

**Function** Sequence-specific DNA-binding transcription factor. Binds to two specific DNA sites located in the

promoter region of HOXA4.

E3 SUMO-protein ligase helping SUMO1 conjugation to its coregulators NAB1 and NAB2, whose

sumoylation down-regulates EGR2 own transcriptional activity. [UniProt]