

ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes PSMA5 / Proteasome 19S S5A
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PSMA5 / Proteasome 19S S5A
Species	Human
Immunogen	Recombinant protein fragment corresponding to PSMA5 / Proteasome 19S S5A protein.
Conjugation	Un-conjugated
Alternate Names	Proteasome subunit alpha type-5; Macropain zeta chain; PSC5; Multicatalytic endopeptidase complex zeta chain; EC 3.4.25.1; ZETA; Proteasome zeta chain

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	26 kDa	

Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	PSMA5
Gene Full Name	proteasome subunit alpha 5
Background	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been found for this gene. [provided by RefSeq, Dec 2010]
Function	The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. [UniProt]
Calculated Mw	26 kDa
PTM	Acetylation; Glycoprotein; Phosphoprotein
Cellular Localization	Cytoplasm. Nucleus. [UniProt]

Images




ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody IHC-P image

Immunohistochemistry: Human colon carcinoma stained with ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody at 1:50 dilution.




ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody ICC/IF image

Immunofluorescence: HeLa stained with ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody at 1:50 dilution.




ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody WB image

Western blot: HeLa, 293T and Jurkat stained with ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody at 1:1000 dilution.




ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody IHC-P image

Immunohistochemistry: Rat heart stained with ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody at 1:50 dilution.



ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody WB image

Western blot: Rat brain and Rat spleen stained with ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody at 1:1000 dilution.



ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody IHC-P image

Immunohistochemistry: Mouse spinal cord stained with ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody at 1:50 dilution.

ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody WB image

Western blot: Mouse liver and Mouse brain stained with ARG43646 anti-PSMA5 / Proteasome 19S S5A antibody at 1:1000 dilution.

