

ARG43665
anti-E2F6 antibodyPackage: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes E2F6
Tested Reactivity	Hu, Ms
Tested Application	ChIP, ICC/IF, IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	E2F6
Species	Human
Immunogen	Recombinant protein of Human E2F6
Conjugation	Un-conjugated
Alternate Names	E2F-6; Transcription factor E2F6

Application Instructions

Application table	Application	Dilution
	ChIP	1:50 - 1:200
	ICC/IF	1:50 - 1:200
	IP	1:20 - 1:50
	WB	1:500 - 1:2000

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Observed Size ~32 - 38 kDa

Properties


Form	Liquid
Purification	Affinity purified.
Buffer	PBS, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Concentration	Batch dependent
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	E2F6
Gene Full Name	E2F transcription factor 6
Background	This gene encodes a member of a family of transcription factors that play a crucial role in the control of the cell cycle. The protein encoded by this gene lacks the transactivation and tumor suppressor protein association domains found in other family members, and contains a modular suppression domain that functions in the inhibition of transcription. It interacts in a complex with chromatin modifying factors. There are pseudogenes for this gene on chromosomes 22 and X. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2013]
Function	Inhibitor of E2F-dependent transcription. Binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3'. Has a preference for the 5'-TTTCCCGC-3' E2F recognition site. E2F6 lacks the transcriptional activation and pocket protein binding domains. Appears to regulate a subset of E2F-dependent genes whose products are required for entry into the cell cycle but not for normal cell cycle progression. May silence expression via the recruitment of a chromatin remodeling complex containing histone H3-K9 methyltransferase activity. Overexpression delays the exit of cells from the S-phase. [UniProt]
Research Area	Gene Regulation antibody
Calculated Mw	32 kDa
PTM	Isopeptide bond; Ubl conjugation [UniProt]
Cellular Localization	Nucleus

Images



ARG43665 anti-E2F6 antibody ICC/IF image

Immunofluorescence: U2OS stained with ARG43665 anti-E2F6 antibody.



ARG43665 anti-E2F6 antibody WB image

Western blot: Mouse lung stained with ARG43665 anti-E2F6 antibody at 1:1000 dilution.