

ARG43822 anti-VDAC1 / Porin antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes VDAC1 / Porin
Tested Reactivity	Hu, Ms, Rat
Tested Application	IHC-P, WB
Specificity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	VDAC1
Species	Human
Immunogen	Synthetic peptide of human VDAC1.
Conjugation	Un-conjugated
Alternate Names	VDAC1; Voltage Dependent Anion Channel 1; PORIN; Voltage-Dependent Anion-Selective Channel Protein 1; Outer Mitochondrial Membrane Protein Porin 1; Plasmalemmal Porin; Porin 31HL; Porin 31HM; MGC111064; VDAC-1; Sperm Binding Protein 1a; HVDAC1; VDAC

Application Instructions

Application table	Application	Dilution
	IHC-P	1:50-1:100
	WB	1:500-1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	31 kDa	

Properties

Form	Liquid
Purification	Affinity Purified
Buffer	Tris-Glycine, 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Preservative	0.01% Sodium azide
Stabilizer	40% Glycerol and 0.05% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	VDAC1
Gene Full Name	voltage-dependent anion channel 1
Background	This gene encodes a voltage-dependent anion channel protein that is a major component of the outer mitochondrial membrane. The encoded protein facilitates the exchange of metabolites and ions across the outer mitochondrial membrane and may regulate mitochondrial functions. This protein also forms channels in the plasma membrane and may be involved in transmembrane electron transport. Alternate splicing results in multiple transcript variants. Multiple pseudogenes of this gene are found on chromosomes 1, 2 3, 6, 9, 12, X and Y.[provided by RefSeq, Sep 2010]
Function	Forms a channel through the mitochondrial outer membrane and also the plasma membrane. The channel at the outer mitochondrial membrane allows diffusion of small hydrophilic molecules; in the plasma membrane it is involved in cell volume regulation and apoptosis. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV. The open state has a weak anion selectivity whereas the closed state is cation-selective. [UniProt]
Research Area	Controls and Markers antibody; Metabolism antibody; Signaling Transduction antibody
Calculated Mw	31 kDa
PTM	Acetylation, Isopeptide bond, Phosphoprotein, Ubl conjugation. [UniProt]
Cellular Localization	Cell membrane, Membrane, Mitochondrion, Mitochondrion outer membrane. [UniProt]

Images



ARG43822 anti-VDAC1 / Porin antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human Colon Cancer tissue stained with ARG43822 anti-VDAC1 / Porin antibody at 1:50 dilution.



ARG43822 anti-VDAC1 / Porin antibody WB image

Western blot: Rat kidney and Mouse kidney stained with ARG43822 anti-VDAC1 / Porin antibody at 1:1000 dilution.