

### Product datasheet

info@arigobio.com

# ARG44056 anti-CD172a / SIRP alpha antibody [15-414] (APC)

Package: 100 tests Store at: 4°C

### **Summary**

Product Description APC-conjugated Mouse Monoclonal antibody recognizes CD172a / SIRP alpha.

Tested Reactivity Hu
Tested Application FACS

Host Mouse

**Clonality** Monoclonal

Clone 15-414

Isotype IgG2a

Target Name CD172a / SIRP alpha

Species Human

Immunogen Human CD172a / SIRP alpha fusion protein.

Conjugation APC

Alternate Names SIRPA; Signal Regulatory Protein Alpha; SHPS1; SIRP; BIT; P84; MFR; SHPS-1; PTPNS1; MYD-1; Tyrosine-

Protein Phosphatase Non-Receptor Type Substrate 1; CD172 Antigen-Like Family Member A; Inhibitory Receptor SHPS-1; Macrophage Fusion Receptor; SIRP-ALPHA-1; SIRPalpha2; SIRPalpha2; CD172a; Brain-Immunoglobulin-Like Molecule With Tyrosine-Based Activation Motifs; Brain Ig-Like Molecule With Tyrosine-Based Activation Motifs; Protein Tyrosine Phosphatase, Non-Receptor Type Substrate 1; Tyrosine Phosphatase SHP Substrate 1; Signal-Regulatory Protein Alpha-1; Signal-Regulatory Protein Alpha-2; Signal-Regulatory Protein Alpha-3; Signal-Regulatory Protein Alpha-3; Signal-Regulatory Protein Alpha-3; CD172a Antigen; Myd-1 Antigen; MyD-1 Antigen; Sirp-Alpha-1; Sirp-Alpha-2; Sirp-Alpha-3; CD172A; MYD1; Bit

### **Application Instructions**

Application table	Application	Dilution
	FACS	10 μl / 10^6 cells
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	

## should be determined by the scientist.

### **Properties**

Form Liquid

Purification The purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions.

The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No

reconstitution is necessary.

Buffer PBS (pH 7.4) and 15 mM Sodium azide

Preservative 15 mM Sodium azide

Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

www.arigobio.com argo.nuts about antibodies 1/2

For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol

SIRPA

Gene Full Name

Signal Regulatory Protein Alpha

Background

The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein can be phosphorylated by tyrosine kinases. The phospho-tyrosine residues of this PTP have been shown to recruit SH2 domain containing tyrosine phosphatases (PTP), and serve as substrates of PTPs. This protein was found to participate in signal transduction mediated by various growth factor receptors. CD47 has been demonstrated to be a ligand for this receptor protein. This gene and its product share very high similarity with several other members of the SIRP family. These related genes are located in close proximity to each other on chromosome 20p13. Multiple alternatively spliced transcript variants have been determined for this gene.

**Function** 

Immunoglobulin-like cell surface receptor for CD47. Acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function.

Calculated Mw

55 kDa


PTM

Disulfide bond, Glycoprotein, Phosphoprotein

Cellular Localization

Membrane

### **Images**



ARG44056 anti-CD172a / SIRP alpha antibody [15-414] (APC) FACS image

Flow Cytometry: Human peripheral whole blood stained with ARG44056 anti-CD172a / SIRP alpha antibody [15-414] (APC) at 10  $\mu l$  / 10^6 cells dilution.