

ARG44256 anti-Sonic Hedgehog antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes Sonic Hedgehog
Tested Reactivity	Hu, Ms
Predict Reactivity	Dog, Rat
Tested Application	FACS, ICC/IF, WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	Sonic Hedgehog
Species	Human
Immunogen	Synthetic peptide around the internal region of Human Sonic Hedgehog (DSEALHPLGMAVK)
Conjugation	Un-conjugated
Alternate Names	HPE3; Sonic hedgehog protein; MCOPCB5; HHG-1; HLP3; HHG1; TPT; TPTPS; SHH; SMMCI

Application Instructions

Application table	Application	Dilution
	FACS	10 µg/µl
	ICC/IF	10 µg/µl
	WB	1-3 µg/µl
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	49 kDa	

Properties

Form	Liquid
Purification	Affinity purified
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed


before use.

Note

For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	SHH
Gene Full Name	sonic hedgehog
Background	<p>This gene encodes a protein that is instrumental in patterning the early embryo. It has been implicated as the key inductive signal in patterning of the ventral neural tube, the anterior-posterior limb axis, and the ventral somites. Of three human proteins showing sequence and functional similarity to the sonic hedgehog protein of <i>Drosophila</i>, this protein is the most similar. The protein is made as a precursor that is autocatalytically cleaved; the N-terminal portion is soluble and contains the signalling activity while the C-terminal portion is involved in precursor processing. More importantly, the C-terminal product covalently attaches a cholesterol moiety to the N-terminal product, restricting the N-terminal product to the cell surface and preventing it from freely diffusing throughout the developing embryo. Defects in this protein or in its signalling pathway are a cause of holoprosencephaly (HPE), a disorder in which the developing forebrain fails to correctly separate into right and left hemispheres. HPE is manifested by facial deformities. It is also thought that mutations in this gene or in its signalling pathway may be responsible for VACTERL syndrome, which is characterized by vertebral defects, anal atresia, tracheoesophageal fistula with esophageal atresia, radial and renal dysplasia, cardiac anomalies, and limb abnormalities. Additionally, mutations in a long range enhancer located approximately 1 megabase upstream of this gene disrupt limb patterning and can result in preaxial polydactyly. [provided by RefSeq, Jul 2008]</p>
Function	<p>Intercellular signal essential for a variety of patterning events during development: signal produced by the notochord that induces ventral cell fate in the neural tube and somites, and the polarizing signal for patterning of the anterior-posterior axis of the developing limb bud. Displays both floor plate- and motor neuron-inducing activity. The threshold concentration of N-product required for motor neuron induction is 5-fold lower than that required for floor plate induction. Activates the transcription of target genes by interacting with its receptor PTCH1 to prevent normal inhibition by PTCH1 on the constitutive signaling activity of SMO (By similarity). [UniProt]</p>
Calculated Mw	50 kDa
PTM	<p>The C-terminal domain displays an autoproteolysis activity and a cholesterol transferase activity. Both activities result in the cleavage of the full-length protein and covalent attachment of a cholesterol moiety to the C-terminal of the newly generated N-terminal fragment (N-product). The N-product is the active species in both local and long-range signaling, whereas the C-product has no signaling activity.</p> <p>Cholesterylation is required for N-product targeting to lipid rafts and multimerization.</p> <p>N-palmitoylation of Cys-24 by HHAT is required for N-product multimerization and full activity. [UniProt]</p>
Cellular Localization	<p>SHH protein N-product: Cell membrane; Lipid-anchor. Note=The dual-lipidated sonic hedgehog protein N-product is firmly tethered to the cell membrane where it forms multimers. Further solubilization and release from the cell surface seem to be achieved through different mechanisms, including the interaction with DISP1 and SCUBE2, movement by lipoprotein particles, transport by cellular extensions called cytonemes or by the proteolytic removal of both terminal lipidated peptides [UniProt]</p>




ARG44256 anti-Sonic Hedgehog antibody ICC/IF image

Immunofluorescence: HeLa stained with ARG44256 anti-Sonic Hedgehog antibody at 10 µg/m dilution.



ARG44256 anti-Sonic Hedgehog antibody WB image

Western blot: Human Liver lysate stained with ARG44256 anti-Sonic Hedgehog antibody at 2 µg/ml dilution.



ARG44256 anti-Sonic Hedgehog antibody FACS image

Flow Cytometry: HeLa stained with ARG44256 anti-Sonic Hedgehog antibody at 10 µg/m dilution.