

Product datasheet

info@arigobio.com

ARG44335 anti-SLIT3 antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal recognizes SLIT3

Tested Reactivity Hu

Predict Reactivity Ms, Rat, Cow, Dog, Gpig, Hrs, Rb

Tested Application WB

Host Rabbit

Clonality Polyclonal

Target Name SLIT3

Species Human

Immunogen Synthetic peptide around the N-terminus of Human SLIT3.

Conjugation Un-conjugated

Alternate Names SLIT3; Slit Guidance Ligand 3; MEGF5; Slit-3; SLIT1; SLIL2; Multiple Epidermal Growth Factor-Like

Domains Protein 5; Multiple EGF-Like Domains Protein 5; Multiple EGF Like Domains 5; Slit Homolog 3

Protein

Application Instructions

Predict Reactivity Note Predicted homology based on immunogen sequence: Mouse: 93%; Rabbit: 93%; Rat: 93%; Cow: 100%;

Dog: 100%; Guinea Pig: 93%; Horse: 100%;

Application table Application Dilution

WB $1-5 \mu g/ml$

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Properties

Form Liquid

Purification Affinity purification with immunogen.

Buffer PBS, 0.09% (w/v) Sodium azide and 2% Sucrose.

Preservative 0.09% (w/v) Sodium azide

Stabilizer 2% Sucrose

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Bioinformation

Gene Symbol SLIT3

Gene Full Name Slit Guidance Ligand 3

Background The protein encoded by this gene is secreted, likely interacting with roundabout homolog receptors to

effect cell migration. Two transcript variants encoding different isoforms have been found for this gene.

Function May act as molecular guidance cue in cellular migration, and function may be mediated by interaction


with roundabout homolog receptors.

Calculated Mw 168 kDa

PTM Disulfide bond, Glycoprotein

Cellular Localization Secreted

Images



ARG44335 anti-SLIT3 antibody WB image

Western blot: HepG2, 293T and HeLa stained with ARG44335 anti-SLIT3 antibody at 3 μ g/ml dilution.