

ARG44749 anti-TANK / TRAF2 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody recognizes TANK / TRAF2
Tested Reactivity	Hu
Tested Application	IP, WB
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Target Name	TANK / TRAF2
Species	Human
Conjugation	Un-conjugated
Alternate Names	TRAF-interacting protein; TRAF2; I-TRAF; TRAF family member-associated NF-kappa-B activator; ITRAF

Application Instructions

Application table	Application	Dilution
	IP	10 µg/mL
	WB	1 µg/mL
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Protein A purification
Buffer	PBS with 0.09% sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	TANK
Gene Full Name	TRAF family member-associated NFKB activator
Background	Members of TRAF family associate with activated TNF-R and CD40 and have been implicated to induce

nuclear translocation of NFkB. TANK can bind to all three TRAFs and synergizing with TRAF2 to activate the NFkB signal cascade.

Images

ARG44749 anti-TANK antibody WB image

Western blot: Raji, Ramos, HT-29 and HCT116 stained with ARG44749 anti-TANK antibody at 1 µg/mL dilution.



ARG44749 anti-TANK antibody IP image

Immunoprecipitation: THP1 lysate immunoprecipitated with 2.5 µg of ARG44749 anti-TANK antibody.

