

ARG51263 anti-Synaptotagmin antibody

Package: 100 μl, 50 μl Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Synaptotagmin
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	lgG
Target Name	Synaptotagmin
Species	Human
Immunogen	Peptide sequence around aa.307~311 (G-L-S-D-P) derived from Human Synaptotagmin 1.
Conjugation	Un-conjugated
Alternate Names	P65; Synaptotagmin I; SytI; SVP65; SYT; p65; Synaptotagmin-1

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:100 - 1:200
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recomme should be determined by the scie	ended starting dilutions and the optimal dilutions or concentrations entist.

Properties

Form	Liquid
Purification	Antibodies were produced by immunizing rabbits with KLH-conjugated synthetic peptide. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Buffer	PBS (without Mg2+ and Ca2+, pH 7.4), 150mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol Gene Full Name Background	SYT1 synaptotagmin I The synaptotagmins are integral membrane proteins of synaptic vesicles thought to serve as Ca(2+) sensors in the process of vesicular trafficking and exocytosis. Calcium binding to synaptotagmin I participates in triggering neurotransmitter release at the synapse
Function	May have a regulatory role in the membrane interactions during trafficking of synaptic vesicles at the active zone of the synapse. It binds acidic phospholipids with a specificity that requires the presence of both an acidic head group and a diacyl backbone. A Ca(2+)-dependent interaction between synaptotagmin and putative receptors for activated protein kinase C has also been reported. It can bind to at least three additional proteins in a Ca(2+)-independent manner; these are neurexins, syntaxin and AP2. [UniProt]
Research Area Calculated Mw	Cancer antibody; Metabolism antibody; Neuroscience antibody; Signaling Transduction antibody 48 kDa

Images



ARG51263 anti-Synaptotagmin antibody WB image

Western blot: 20 μg of Mouse brain and Rat brain lysates stained with ARG51263 anti-Synaptotagmin antibody at 1:500 dilution.



ARG51263 anti-Synaptotagmin antibody ICC/IF image

Immunofluorescence: methanol-fixed HeLa cells stained with anti-Synaptotagmin antibody ARG51263.