

## ARG51731 anti-CDK1 / CDC2 phospho (Tyr15) antibody

Package: 100 µl, 50 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes CDK1 / CDC2 phospho (Tyr15)
Tested Reactivity	Hu, Ms, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CDK1 / CDC2
Species	Human
Immunogen	Peptide sequence around phosphorylation site of tyrosine 15 (G-T-Y(p)-G-V) derived from Human CDC2.
Conjugation	Un-conjugated
Alternate Names	CDK1; Cyclin Dependent Kinase 1; CDC28A; CDC2; Cell Division Cycle 2, G1 To S And G2 To M; Cell Division Control Protein 2 Homolog; Cell Division Protein Kinase 1; Cyclin-Dependent Kinase 1; P34 Protein Kinase; P34CDC2; Cell Cycle Controller CDC2; EC 2.7.11.22; EC 2.7.11.23; CDKN1

### Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


### Properties

Form	Liquid
Purification	Antibodies were produced by immunizing rabbits with KLH-conjugated synthetic phosphopeptide. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. In addition, non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Buffer	PBS (without Mg2+ and Ca2+, pH 7.4), 150mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	CDK1
Gene Full Name	cyclin-dependent kinase 1
Background	<p>The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2023]</p>
Function	<p>Plays a key role in the control of the eukaryotic cell cycle by modulating the centrosome cycle as well as mitotic onset; promotes G2-M transition via association with multiple interphase cyclins. [UniProt]</p>
Research Area	Cell Biology and Cellular Response antibody; Gene Regulation antibody; Neuroscience antibody
Calculated Mw	34 kDa
PTM	<p>Phosphorylation at Thr-14 and Tyr-15 by PKMYT1 prevents nuclear translocation.</p> <p>Phosphorylation at Tyr-15 by WEE1 and WEE2 inhibits the protein kinase activity and acts as a negative regulator of entry into mitosis (G2 to M transition).</p> <p>Dephosphorylation by active CDC25A and CDC25B at Thr-14 and Tyr-15, leads to CDK1 activation at the G2-M transition.</p> <p>Phosphorylation at Tyr-15 by WEE2 during oogenesis is required to maintain meiotic arrest in oocytes during the germinal vesicle (GV) stage, a long period of quiescence at dictyate prophase I, leading to prevent meiotic reentry.</p> <p>In response to UV irradiation, phosphorylation at Tyr-15 by PRKCD activates the G2/M DNA damage checkpoint. [UniProt]</p>
Cellular Localization	Cytoplasm, Cytoskeleton, Mitochondrion, Nucleus. [UniProt]

Images




ARG51731 anti-CDK1 / CDC2 phospho (Tyr15) antibody WB image

Western blot: Extracts from HeLa cells untreated(lane 1) or treated with UV(lane 2) stained with ARG51731 anti-CDK1 / CDC2 phospho (Tyr15) antibody.



ARG51731 anti-CDK1 / CDC2 phospho (Tyr15) antibody WB image

Western blot: Extracts from 293 cells, treated with Hydroxyurea or calf intestinal phosphatase (CIP), stained with ARG51731 anti-CDK1 / CDC2 phospho (Tyr15) antibody.



ARG51731 anti-CDK1 / CDC2 phospho (Tyr15) antibody WB image

Western blot: Extracts from 293 cells, treated with Hydroxyurea or calf intestinal phosphatase (CIP), stained with ARG51731 anti-CDK1 / CDC2 phospho (Tyr15) antibody.