

Product datasheet

info@arigobio.com

ARG52431 anti-Synapsin 1 phospho (Ser549) antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes Synapsin 1 phospho (Ser549)

Tested Reactivity Rat

Predict Reactivity Hu, Ms, Bov, Dog, NHuPrm

Tested Application IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name Synapsin 1

Species Rat

Immunogen KLH-conjugated phosphospecific peptide around Ser549 of Rat Synapsin 1.

Conjugation Un-conjugated

Alternate Names SYNI; Brain protein 4.1; Synapsin-1; SYN1a; SYN1b; Synapsin I

Application Instructions

Application table	Application	Dilution
	IHC-P	1:500
	WB	1:1000
Application Note	Specific for $^{\sim}78k$ synapsin I doublet phosphorylated at Ser549. Immunolabeling of the synapsin I band is blocked by λ -phosphatase treatment. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Note

Form	Liquid	
Purification	Affinity Purified	
Buffer	10 mM HEPES (pH 7.5), 150 mM NaCl, 0.1 mg/ml BSA and 50% Glycerol	
Stabilizer	0.1 mg/ml BSA, 50% Glycerol	
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.	

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GenelD: 24949 Rat

Swiss-port # P09951 Rat

Gene Symbol Syn1

Gene Full Name synapsin I

Background Synapsin I plays a key role in synaptic plasticity in brain (Feng et al., 2002; Nayak et al., 1996). This

effect is due in large part to the ability of the synapsins to regulate the availability of synaptic vesicles

for release. The role of synapsin in synaptic plasticity and in synaptogensis is regulated by

phosphorylation (Jovanovic et al., 2001; Kao et al., 2002). Ser 549 along with Ser 62 and Ser 67 are the sites of synapsin I that are phosphorylated by MAP kinase (Jovanovic et al., 1996). Phosphorylation and subsequent dephosphorylation of this site is thought to play a key role in synaptic vesicle trafficking.

Function Neuronal phosphoprotein that coats synaptic vesicles, binds to the cytoskeleton, and is believed to

function in the regulation of neurotransmitter release. [UniProt]

Research Area Neuroscience antibody


Calculated Mw 74 kDa

PTM Substrate of at least four different protein kinases. It is probable that phosphorylation plays a role in

the regulation of synapsin-1 in the nerve terminal.

Phosphorylation at Ser-9 dissociates synapsins from synaptic vesicles.

Images



ARG52431 anti-Synapsin 1 phospho (Ser549) antibody WB image

Western blot: Rat cortex lysate showing specific immunolabeling of the ~78 kDa Synapsin 1 phosphorylated at Ser549 (Control) stained with ARG52431 anti-Synapsin 1 phospho (Ser549) antibody. Phosphospecificity is shown in the second lane (lambda-phosphatase: λ -Ptase).