

ARG53467 anti-PIK3CA / p110 alpha antibody [SP139]

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Monoclonal antibody [SP139] recognizes PIK3CA / p110 alpha
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Monoclonal
Clone	SP139
Isotype	IgG
Target Name	PIK3CA / p110 alpha
Species	Human
Immunogen	Synthetic peptide in the internal region of the human p13KCA protein.
Conjugation	Un-conjugated
Alternate Names	MCM; MCMTC; p110alpha; PI3-kinase subunit alpha; PI3K; CWS5; p110-alpha; PI3Kalpha; CLOVE; EC 2.7.1.153; MCAP; PtdIns-3-kinase subunit p110-alpha; EC 2.7.11.1; Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform; Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit alpha; PI3K-alpha; PtdIns-3-kinase subunit alpha; Phosphoinositide-3-kinase catalytic alpha polypeptide; Serine/threonine protein kinase PIK3CA

Application Instructions

Application table	Application	Dilution
	IHC-P	1:100
Application Note	IHC-P: Antigen Retrieval: Boil tissue section in EDTA buffer, pH 8.0 for 10 min followed by cooling at room temperature for 20 min. Incubation Time: 10 min at RT. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Breast Carcinoma	

Properties

Form	Liquid
Purification	Purified by protein A/G
Buffer	PBS (pH 7.6), 1% BSA and < 0.1% Sodium azide
Preservative	< 0.1% Sodium azide
Stabilizer	1% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot


and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 5290 Human Swiss-port # P42336 Human
Gene Symbol	PIK3CA
Gene Full Name	phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha
Background	Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. [provided by RefSeq, Jul 2008]
Function	Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns(4,5)P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors. Involved in the activation of AKT1 upon stimulation by receptor tyrosine kinases ligands such as EGF, insulin, IGF1, VEGFA and PDGF. Involved in signaling via insulin-receptor substrate (IRS) proteins. Essential in endothelial cell migration during vascular development through VEGFA signaling, possibly by regulating RhoA activity. Required for lymphatic vasculature development, possibly by binding to RAS and by activation by EGF and FGF2, but not by PDGF. Regulates invadopodia formation in breast cancer cells through the PDK1-AKT1 pathway. Participates in cardiomyogenesis in embryonic stem cells through a AKT1 pathway. Participates in vasculogenesis in embryonic stem cells through PDK1 and protein kinase C pathway. Has also serine-protein kinase activity: phosphorylates PIK3R1 (p85alpha regulatory subunit), EIF4EBP1 and HRAS. [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Metabolism antibody; Neuroscience antibody
Calculated Mw	124 kDa
Cellular Localization	Membrane

Images




ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Breast Ductal Carcinoma stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].




ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Colon Adenocarcinoma stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].




ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Stomach stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].




ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Stomach Adenocarcinoma stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].




ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Bladder Transitional Cell Carcinoma stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].




ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Prostate Adenocarcinoma stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].



ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Skin Squamous Cell Carcinoma stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].



ARG53467 anti-PIK3CA / p110 alpha antibody [SP139] IHC-P image

Immunohistochemistry: Human Skin stained with ARG53467 anti-PIK3CA / p110 alpha antibody [SP139].
