

## ARG53908 anti-CD86 antibody [BU63] (PE)

Package: 100 tests  
Store at: 4°C

### Summary

Product Description	PE-conjugated Mouse Monoclonal antibody [BU63] recognizes CD86
Tested Reactivity	Hu
Tested Application	FACS
Specificity	The clone BU63 reacts with CD86 (B7-2), a 70 kDa type I transmembrane glycoprotein of immunoglobulin supergene family, expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes. HLDA V; WS Code BP BP072 HLDA V; WS Code A A109 HLDA VI; WS Code BP 95 HLDA VI; WS Code B CD86.9
Host	Mouse
Clonality	Monoclonal
Clone	BU63
Isotype	IgG1
Target Name	CD86
Immunogen	B-lymphoblastoid cell line ARH 77
Conjugation	PE
Alternate Names	B70; B7.2; LAB72; CD antigen CD86; B7-2; FUN-1; CD28LG2; T-lymphocyte activation antigen CD86; CTLA-4 counter-receptor B7.2; Activation B7-2 antigen; BU63

### Application Instructions

Application table	Application	Dilution
	FACS	20 µl / 10 <sup>6</sup> cells

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Purification Note	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Buffer	PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
Preservative	15 mM Sodium azide
Stabilizer	0.2% (w/v) high-grade protease free BSA

**Storage instruction** Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


**Note** For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 942 Human</a> <a href="#">Swiss-port # P42081 Human</a>
Gene Symbol	CD86
Gene Full Name	CD86 molecule
Background	CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, the both molecules can also substitute for each other in this process. The question what are the differences in CD80 and CD86 competency has not been fully elucidated yet; there are still conflicts in results about their respective roles in initiation or sustaining of the T cell immune response.
Function	Receptor involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. Isoform 2 interferes with the formation of CD86 clusters, and thus acts as a negative regulator of T-cell activation. [UniProt]
Research Area	Developmental Biology antibody; Immune System antibody; Microbiology and Infectious Disease antibody
Calculated Mw	38 kDa
PTM	Polyubiquitinated; which is promoted by MARCH8 and results in endocytosis and lysosomal degradation.

## Images



ARG53908 anti-CD86 antibody [BU63] (PE) FACS image

Flow Cytometry: Human peripheral blood cells stained with ARG53908 anti-CD86 antibody [BU63] (PE).