

## ARG54039 anti-CD54 / ICAM1 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody recognizes CD54 / ICAM1
Tested Reactivity	Hu
Tested Application	ICC/IF, WB
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Target Name	CD54 / ICAM1
Species	Human
Immunogen	Purified recombinant human CD54 / ICAM1 protein fragments expressed in E.coli.
Conjugation	Un-conjugated
Alternate Names	CD54; CD antigen CD54; BB2; P3.58; Intercellular adhesion molecule 1; Major group rhinovirus receptor; ICAM-1

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:200
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	96 kDa	


### Properties

Form	Liquid
Purification	Affinity purified
Buffer	PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Concentration	2.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	<a href="#">GeneID: 3383 Human</a> <a href="#">Swiss-port # P05362 Human</a>
Gene Symbol	ICAM1
Gene Full Name	intercellular adhesion molecule 1
Background	This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor. [provided by RefSeq, Jul 2008]
Function	ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation (By similarity). [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Developmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibody
Calculated Mw	58 kDa
PTM	Monoubiquitinated, which is promoted by MARCH9 and leads to endocytosis.
Cellular Localization	Membrane; Single-pass type I membrane protein.

Images



ARG54039 anti-CD54 / ICAM1 antibody ICC/IF image

Immunofluorescence: SW480 cells fixed by anhydrous methanol for 2 hours at -20°C and stained with ARG54039 anti-CD54 / ICAM1 antibody at 1:200 dilution.



Raji

ARG54039 anti-CD54 / ICAM1 antibody WB image

Western blot: Raji cell lysate stained with ARG54039 anti-CD54 / ICAM1 antibody at 1:1000 dilution.