

## ARG54156 anti-Hsp 90 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody recognizes Hsp 90
Tested Reactivity	Hu, Ms, Rat
Tested Application	WB
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Target Name	Hsp 90
Species	Human
Immunogen	Purified recombinant human HSP90AA1 protein fragments expressed in E.coli
Conjugation	Un-conjugated
Alternate Names	EL52; Hsp90; HSPC1; Heat shock 86 kDa; LAP-2; HSP90N; LAP2; HSP90A; HSPCAL4; HSP89A; HSP86; HSP 86; HSPCA; Lipopolysaccharide-associated protein 2; HSPCAL1; LPS-associated protein 2; HSPN; Renal carcinoma antigen NY-REN-38; Heat shock protein HSP 90-alpha; Hsp89

### Application Instructions

Application table	Application	Dilution
	WB	1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	90 kDa	


### Properties

Form	Liquid
Purification	Affinity purified
Buffer	PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Concentration	2.25 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Gene Symbol	HSP90AA1
Gene Full Name	heat shock protein 90kDa alpha (cytosolic), class A member 1
Background	Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function.
Function	Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function. Binds bacterial lipopolysaccharide (LPS) et mediates LPS-induced inflammatory response, including TNF secretion by monocytes. [UniProt]
Research Area	Cancer antibody; Signaling Transduction antibody
Calculated Mw	85 kDa
PTM	ISGylated. S-nitrosylated; negatively regulates the ATPase activity and the activation of eNOS by HSP90AA1.
Cellular Localization	Cytoplasm. Melanosome. Note: Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

## Images



ARG54156 anti-Hsp 90 antibody WB image

Western blot: 3T3 cell lysate stained with ARG54156 anti-Hsp 90 antibody at 1:2000 dilution.