

## ARG54373 anti-Survivin antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes Survivin
Tested Reactivity	Ms
Tested Application	IHC-P, WB
Specificity	This antibody recognizes mouse Survivin (17kDa).
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Survivin
Species	Mouse
Immunogen	Synthetic peptide corresponding to aa 128-140 of mouse Survivin (accession no. BAA28266).
Conjugation	Un-conjugated
Alternate Names	API4; Apoptosis inhibitor 4; EPR-1; Apoptosis inhibitor survivin; Baculoviral IAP repeat-containing protein 5

### Application Instructions


Application table	Application	Dilution
	IHC-P	10-20 µg/mL
	WB	1 µg/mL
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse spleen	

### Properties

Form	Liquid
Purification	Immunoaffinity chroma-tography
Buffer	PBS (pH 7.4) and 0.02% Sodium azide
Preservative	0.02% Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Database links	<a href="#">GeneID: 11799 Mouse</a> <a href="#">Swiss-port # O70201 Mouse</a>
Gene Symbol	Birc5
Gene Full Name	baculoviral IAP repeat-containing 5
Background	Apoptosis can be prevented by the inhibitor of apoptosis proteins (IAPs). IAPs constitute a conserved gene family that binds to and inhibits cell death proteases. A novel IAP has been identified and designated Survivin. Survivin interacts with the processed form of caspase-3 and inhibits its proteolytic activity. Survivin is expressed predominantly in tissues of embryos, transformed cell lines, and many human cancers.
Function	Multitasking protein that has dual roles in promoting cell proliferation and preventing apoptosis. Component of a chromosome passage protein complex (CPC) which is essential for chromosome alignment and segregation during mitosis and cytokinesis. Acts as an important regulator of the localization of this complex; directs CPC movement to different locations from the inner centromere during prometaphase to midbody during cytokinesis and participates in the organization of the center spindle by associating with polymerized microtubules. The complex with RAN plays a role in mitotic spindle formation by serving as a physical scaffold to help deliver the RAN effector molecule TPX2 to microtubules. May counteract a default induction of apoptosis in G2/M phase. The acetylated form represses STAT3 transactivation of target gene promoters. May play a role in neoplasia. Inhibitor of CASP3 and CASP7 (By similarity). [UniProt]
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Cell Death antibody; Neuroscience antibody
Calculated Mw	16 kDa
PTM	Ubiquitinated by the Cul9-RING ubiquitin-protein ligase complex, leading to its degradation. Ubiquitination is required for centrosomal targeting. In vitro phosphorylation at Thr-117 by AURKB prevents interaction with INCENP and localization to mitotic chromosomes (PubMed:14610074). Phosphorylation at Thr-48 by CK2 is critical for its mitotic and anti-apoptotic activities (PubMed:21252625). Phosphorylation at Thr-34 by CDK15 is critical for its anti-apoptotic activity (PubMed:24866247). Phosphorylation at Ser-20 by AURKC is critical for regulation of proper chromosome alignment and segregation, and possibly cytokinesis. Acetylation at Lys-129 by CBP results in its homodimerization, while deacetylation promotes the formation of monomers which heterodimerize with XPO1/CRM1 which facilitates its nuclear export. The acetylated form represses STAT3 transactivation. The dynamic equilibrium between its acetylation and deacetylation at Lys-129 determines its interaction with XPO1/CRM1, its subsequent subcellular localization, and its ability to inhibit STAT3 transactivation.

## Images



ARG54373 anti-Survivin antibody WB image

Western Blot: mouse spleen stained with ARG54373 anti-Survivin antibody at 1 µg/ml dilution.



ARG54373 anti-Survivin antibody IHC-P image

Immunohistochemistry: mouse spleen stained with ARG54373 anti-Survivin antibody at 10 µg/ml dilution.