

ARG54403 anti-Daxx antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Daxx
Tested Reactivity	Hu
Tested Application	ICC/IF, WB
Specificity	This antibody recognizes full-length human, monkey, and mouse Daxx (120kDa).
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Daxx
Species	Human
Immunogen	Peptide corresponding to aa 722-740 at the C-terminus of human Daxx (accession no. CAG33366).
Conjugation	Un-conjugated
Alternate Names	Daxx; ETS1-associated protein 1; hDaxx; Fas death domain-associated protein; Death domain-associated protein 6; DAP6; EAP1; BING2


Application Instructions

Application table	Application	Dilution
	ICC/IF	Assay-dependent
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	293 and HeLa	

Properties


Form	Liquid
Purification	Immunoaffinity chroma-tography
Buffer	PBS (pH 7.4) and 0.02% Sodium azide
Preservative	0.02% Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Database links	GeneID: 1616 Human Swiss-port # Q9UER7 Human
Gene Symbol	DAXX
Gene Full Name	death-domain associated protein
Background	A novel Fas binding protein was recently cloned from human and mouse and designated Daxx. Daxx binds specifically to the Fas death domain, enhances Fas induced apoptosis, and activates the Jun N-terminal kinase (JNK) pathway. Daxx is widely expressed in fetal and adult human and mouse tissues, indicating its important function in Fas signaling pathways.
Function	<p>Transcription corepressor known to repress transcriptional potential of several sumoylated transcription factors. Down-regulates basal and activated transcription. Its transcription repressor activity is modulated by recruiting it to subnuclear compartments like the nucleolus or PML/POD/ND10 nuclear bodies through interactions with MCSR1 and PML, respectively. Seems to regulate transcription in PML/POD/ND10 nuclear bodies together with PML and may influence TNFRSF6-dependent apoptosis thereby. Inhibits transcriptional activation of PAX3 and ETS1 through direct protein-protein interactions. Modulates PAX5 activity; the function seems to involve CREBBP. Acts as an adapter protein in a MDM2-DAXX-USP7 complex by regulating the RING-finger E3 ligase MDM2 ubiquitination activity. Under non-stress condition, in association with the deubiquitinating USP7, prevents MDM2 self-ubiquitination and enhances the intrinsic E3 ligase activity of MDM2 towards TP53, thereby promoting TP53 ubiquitination and subsequent proteasomal degradation. Upon DNA damage, its association with MDM2 and USP7 is disrupted, resulting in increased MDM2 autoubiquitination and consequently, MDM2 degradation, which leads to TP53 stabilization. Acts as histone chaperone that facilitates deposition of histone H3.3. Acts as targeting component of the chromatin remodeling complex ATRX:DAXX which has ATP-dependent DNA translocase activity and catalyzes the replication-independent deposition of histone H3.3 in pericentric DNA repeats outside S-phase and telomeres, and the in vitro remodeling of H3.3-containing nucleosomes. Does not affect the ATPase activity of ATRX but alleviates its transcription repression activity. Upon neuronal activation associates with regulatory elements of selected immediate early genes where it promotes deposition of histone H3.3 which may be linked to transcriptional induction of these genes. Required for the recruitment of histone H3.3:H4 dimers to PML-nuclear bodies (PML-NBs); the process is independent of ATRX and facilitated by ASF1A; PML-NBs are suggested to function as regulatory sites for the incorporation of newly synthesized histone H3.3 into chromatin. In case of overexpression of centromeric histone variant CENPA (as found in various tumors) is involved in its mislocalization to chromosomes; the ectopic localization involves a heterotypic tetramer containing CENPA, and histones H3.3 and H4 and decreases binding of CTCF to chromatin. Proposed to mediate activation of the JNK pathway and apoptosis via MAP3K5 in response to signaling from TNFRSF6 and TGFBR2. Interaction with HSPB1/HSP27 may prevent interaction with TNFRSF6 and MAP3K5 and block DAXX-mediated apoptosis. In contrast, in lymphoid cells JNK activation and TNFRSF6-mediated apoptosis may not involve DAXX. Shows restriction activity towards human cytomegalovirus (HCMV). [UniProt]</p>
Research Area	Cell Biology and Cellular Response antibody; Cell Death antibody; Gene Regulation antibody; Microbiology and Infectious Disease antibody
Calculated Mw	81 kDa
PTM	<p>Sumoylated with SUMO1 on multiple lysine residues.</p> <p>Phosphorylated by HIPK1 upon glucose deprivation.</p> <p>Polyubiquitinated; which is promoted by CUL3 and SPOP and results in proteasomal degradation.</p> <p>Ubiquitinated by MDM2; inducing its degradation. Deubiquitinated by USP7; leading to stabilize it.</p>



ARG54403 anti-Daxx antibody WB image

Western Blot: 293 stained with ARG54403 anti-Daxx antibody at 1 μ g/ml dilution.



ARG54403 anti-Daxx antibody ICC/IF image

Immunofluorescence: HeLa stained with ARG54403 anti-Daxx antibody at 10 μ g/ml dilution.