

## ARG55300 anti-CES1 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes CES1
Tested Reactivity	Hu, Ms
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CES1
Species	Human
Immunogen	Recombinant protein of Human CES1
Conjugation	Un-conjugated
Alternate Names	SES1; Cocaine carboxylesterase; REH; ACAT; Serine esterase 1; HMSE; Retinyl ester hydrolase; Methylumbelliferyl-acetate deacetylase 1; hCE-1; EC 3.1.1.56; CE-1; CEH; HMSE1; TGH; PCE-1; Acyl-coenzyme A:cholesterol acyltransferase; EC 3.1.1.1; Liver carboxylesterase 1; Brain carboxylesterase hBr1; Triacylglycerol hydrolase; CES2; Egasyn; Carboxylesterase 1; Monocyte/macrophage serine esterase

### Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A549 and Mouse liver	


### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links	<a href="#">GeneID: 1066 Human</a> <a href="#">Swiss-port # P23141 Human</a>
Gene Symbol	CES1
Gene Full Name	carboxylesterase 1
Background	This gene encodes a member of the carboxylesterase large family. The family members are responsible for the hydrolysis or transesterification of various xenobiotics, such as cocaine and heroin, and endogenous substrates with ester, thioester, or amide bonds. They may participate in fatty acyl and cholesterol ester metabolism, and may play a role in the blood-brain barrier system. This enzyme is the major liver enzyme and functions in liver drug clearance. Mutations of this gene cause carboxylesterase 1 deficiency. Three transcript variants encoding three different isoforms have been found for this gene. [provided by RefSeq, Jun 2010]
Function	Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Hydrolyzes aromatic and aliphatic esters, but has no catalytic activity toward amides or a fatty acyl-CoA ester. Hydrolyzes the methyl ester group of cocaine to form benzoylecgonine. Catalyzes the transesterification of cocaine to form cocaethylene. Displays fatty acid ethyl ester synthase activity, catalyzing the ethyl esterification of oleic acid to ethyloleate. [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Metabolism antibody; Signaling Transduction antibody
Calculated Mw	63 kDa
PTM	Contains sialic acid. Cleavage of the signal sequence can occur at 2 positions, either between Trp-17 and Gly-18 or between Gly-18 and His-19.

## Images



ARG55300 anti-CES1 antibody WB image

Western blot: A549 and Mouse liver lysates stained with ARG55300 anti-CES1 antibody.