

ARG55353 anti-ALDH1A1 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody recognizes ALDH1A1	
Tested Reactivity	Hu	
Tested Application	ICC/IF, IHC-P, WB	
Host	Mouse	
Clonality	Monoclonal	
Clone	152CT1.2.2	
Isotype	lgG1	
Target Name	ALDH1A1	
Species	Human	
Immunogen	Human ALDH1A1 recombinant protein.	
Conjugation	Un-conjugated	
Alternate Names	HEL-S-53e; RalDH1; Aldehyde dehydrogenase family 1 member A1; Retinal dehydrogenase 1; HEL-9; RALDH 1; ALDH1; Aldehyde dehydrogenase, cytosolic; ALDH11; EC 1.2.1.36; RALDH1; HEL12; PUMB1; ALHDII; ALDC; ALDH-E1	

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:10 - 1:50
	IHC-P	1:50 - 1:100
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	K562	


Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% (W/V) Sodium azide
Preservative	0.09% (W/V) Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

		1.1.2.2
RIUIN	forma	tion
DIOIII	101111u	CIOIL


Database links	GenelD: 216 Human
	Swiss-port # P00352 Human
Gene Symbol	ALDH1A1
Gene Full Name	aldehyde dehydrogenase 1 family, member A1
Background	The protein encoded by this gene belongs to the aldehyde dehydrogenase family. Aldehyde dehydrogenase is the next enzyme after alcohol dehydrogenase in the major pathway of alcohol metabolism. There are two major aldehyde dehydrogenase isozymes in the liver, cytosolic and mitochondrial, which are encoded by distinct genes, and can be distinguished by their electrophoretic mobility, kinetic properties, and subcellular localization. This gene encodes the cytosolic isozyme. Studies in mice show that through its role in retinol metabolism, this gene may also be involved in the regulation of the metabolic responses to high-fat diet. [provided by RefSeq, Mar 2011]
Function	Binds free retinal and cellular retinol-binding protein-bound retinal. Can convert/oxidize retinaldehyde to retinoic acid (By similarity). [UniProt]
Research Area	Cancer antibody; Developmental Biology antibody; Neuroscience antibody
Calculated Mw	55 kDa
PTM	The N-terminus is blocked most probably by acetylation.
Cellular Localization	Cytoplasm.

Images




ARG55353 anti-ALDH1A1 antibody ICC/IF image

Immunofluorescence: NCI-H460 cells stained with ARG55353 anti-ALDH1A1 antibody (green). DAPI (blue) for nuclear staining.



ARG55353 anti-ALDH1A1 antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human colon carcinoma stained with ARG55353 anti-ALDH1A1 antibody.



ARG55353 anti-ALDH1A1 antibody WB image

Western blot: 20 μg of K562 cell lysate stained with ARG55353 anti-ALDH1A1 antibody at 1:1000 dilution.