

**ARG55394**  
anti-MED8 antibodyPackage: 100 µl  
Store at: -20°C

## Summary

Product Description	Rabbit Polyclonal antibody recognizes MED8
Tested Reactivity	Hu
Predict Reactivity	Ms, Zfsh
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MED8
Species	Human
Immunogen	KLH-conjugated synthetic peptide corresponding to aa. 95-122 (Center) of Human MED8.
Conjugation	Un-conjugated
Alternate Names	Mediator of RNA polymerase II transcription subunit 8; Activator-recruited cofactor 32 kDa component; ARC32; Mediator complex subunit 8

## Application Instructions

Application table	Application	Dilution
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HL-60	


## Properties

Form	Liquid
Purification	Purification with Protein A and immunogen peptide.
Buffer	PBS and 0.09% (W/V) Sodium azide
Preservative	0.09% (W/V) Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links	<a href="#">GeneID: 112950 Human</a> <a href="#">Swiss-port # Q96G25 Human</a>
Gene Symbol	MED8
Gene Full Name	mediator complex subunit 8
Background	This gene encodes a protein component of the mediator complex, which aids in transcriptional activation through interaction with RNA polymerase II and gene-specific transcription factors. The encoded protein may also function in ubiquitin ligation and protein degradation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2013]
Function	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May play a role as a target recruitment subunit in E3 ubiquitin-protein ligase complexes and thus in ubiquitination and subsequent proteasomal degradation of target proteins. [UniProt]
Research Area	Gene Regulation antibody
Calculated Mw	29 kDa

## Images



ARG55394 anti-MED8 antibody WB image

Western blot: 35 µg of HL-60 cell lysate stained with ARG55394 anti-MED8 antibody.