

## ARG55434 anti-CREB3L4 / AIBZIP antibody

Package: 100 µl  
Store at: -20°C

### Summary

|                     |   |
|---------------------|---|
| Product Description | Rabbit Polyclonal antibody recognizes CREB3L4 / AIBZIP  |
| Tested Reactivity   | Hu  |
| Tested Application  | WB  |
| Host                | Rabbit  |
| Clonality           | Polyclonal  |
| Isotype             | IgG   |
| Target Name         | CREB3L4 / AIBZIP  |
| Species             | Human   |
| Immunogen           | KLH-conjugated synthetic peptide corresponding to aa. 338-366 (C-terminus) of Human CREB3L4.  |
| Conjugation         | Un-conjugated   |
| Alternate Names     | Transcript induced in spermiogenesis protein 40; cAMP-responsive element-binding protein 3-like protein 4; Androgen-induced basic leucine zipper protein; Cyclic AMP-responsive element-binding protein 3-like protein 4; cAMP-responsive element-binding protein 4; CREB4; CREB3; hJAL; Tisp40; ATCE1; JAL; Cyclic AMP-responsive element-binding protein 4; Attaching to CRE-like 1; CREB-4; AibZIP; AIBZIP |

### Application Instructions

| Application table | <table> <tr> <th>Application</th><th>Dilution</th></tr> <tr> <td>WB</td><td>1:1000</td></tr> </table>                                      | Application | Dilution | WB | 1:1000 |
|-------------------|--|-------------|----------|----|--------|
| Application       | Dilution   |             |          |    |        |
| WB                | 1:1000   |             |          |    |        |
| Application Note  | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |             |          |    |        |
| Positive Control  | PC-3   |             |          |    |        |


### Properties

|                     |  |
|---------------------|--|
| Form                | Liquid   |
| Purification        | Purification with Protein A and immunogen peptide.   |
| Buffer              | PBS and 0.09% (W/V) Sodium azide   |
| Preservative        | 0.09% (W/V) Sodium azide   |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note                | For laboratory research only, not for drug, diagnostic or other use.   |

## Bioinformation

|                       |  |
|-----------------------|--|
| Database links        | <a href="#">GeneID: 148327 Human</a><br><a href="#">Swiss-port # Q8TEY5 Human</a>  |
| Gene Symbol           | CREB3L4  |
| Gene Full Name        | cAMP responsive element binding protein 3-like 4   |
| Background            | This gene encodes a CREB (cAMP responsive element binding) protein with a transmembrane domain which localizes it to the ER membrane. The encoded protein is a transcriptional activator which contains a dimerization domain, and this protein may function in a number of processing pathways including protein processing. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2011] |
| Function              | Transcriptional activator that may play a role in the unfolded protein response. Binds to the UPR element (UPRE) but not to CRE element. Preferentially binds DNA with to the consensus sequence 5'-T[GT]ACGT[GA][GT]-3' and has transcriptional activation activity from UPRE. Binds to NF-kappa-B site and has transcriptional activation activity from NF-kappa-B-containing regulatory elements (By similarity). [UniProt]                       |
| Research Area         | Gene Regulation antibody   |
| Calculated Mw         | 43 kDa   |
| PTM                   | N-glycosylated in the C-terminal region.<br>Controlled by regulated intramembrane proteolysis (RIP). Following ER stress a fragment containing the cytoplasmic transcription factor domain is released by proteolysis. The cleavage seems to be performed sequentially by site-1 and site-2 proteases (PS1 and PS2). PS1 cleavage may be suppressed by a determinant in the C-terminal region (By similarity).                                       |
| Cellular Localization | Endoplasmic reticulum membrane; Single-pass type II membrane protein. Golgi apparatus membrane; Single-pass type II membrane protein. Note=May also be located in Golgi apparatus  |

## Images



ARG55434 anti-CREB3L4 / AIBZIP antibody WB image

Western blot: 35 µg of PC-3 cell lysate stained with ARG55434 anti-CREB3L4 / AIBZIP antibody.