

ARG55567 anti-Cathepsin D antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody recognizes Cathepsin D
Tested Reactivity	Hu
Tested Application	IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Clone	892CT11.1.1
Isotype	IgG1
Target Name	Cathepsin D
Species	Human
Immunogen	Purified His-tagged Human Cathepsin D protein.
Conjugation	Un-conjugated
Alternate Names	CPSD; EC 3.4.23.5; HEL-S-130P; CLN10; Cathepsin D

Application Instructions

Application table	Application	Dilution
	IHC-P	1:25
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	MCF7	


Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% (W/V) Sodium azide
Preservative	0.09% (W/V) Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 1509 Human Swiss-port # P07339 Human
Gene Symbol	CTSD
Gene Full Name	cathepsin D
Background	This gene encodes a lysosomal aspartyl protease composed of a dimer of disulfide-linked heavy and light chains, both produced from a single protein precursor. This proteinase, which is a member of the peptidase C1 family, has a specificity similar to but narrower than that of pepsin A. Transcription of this gene is initiated from several sites, including one which is a start site for an estrogen-regulated transcript. Mutations in this gene are involved in the pathogenesis of several diseases, including breast cancer and possibly Alzheimer disease. [provided by RefSeq, Jul 2008]
Function	Acid protease active in intracellular protein breakdown. Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease. [UniProt]
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Neuroscience antibody; Signaling Transduction antibody
Calculated Mw	45 kDa
PTM	N- and O-glycosylated. Undergoes proteolytic cleavage and activation by ADAM30. As well as the major heavy chain which starts at Leu-169, 2 minor forms starting at Gly-170 and Gly-171 have been identified (PubMed:1426530). An additional form starting at Ala-168 has also been identified (PubMed:27333034).
Cellular Localization	Lysosome. Melanosome. Secreted, extracellular space. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. In aortic samples, detected as an extracellular protein loosely bound to the matrix (PubMed:20551380).

Images



ARG55567 anti-Cathepsin D antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human liver section stained with ARG55567 anti-Cathepsin D antibody at 1:25 dilution.



ARG55567 anti-Cathepsin D antibody WB image

Western blot: 35 μ g of MCF7 cell lysate stained with ARG55567 anti-Cathepsin D antibody.