

ARG55686 anti-Hsp 60 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Hsp 60
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat, Bov, Chk, Hm
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Hsp 60
Species	Human
Immunogen	KLH-conjugated synthetic peptide corresponding to aa. 187-215 (Center) of Human Hsp60.
Conjugation	Un-conjugated
Alternate Names	Heat shock protein 60; HuCHA60; 60 kDa heat shock protein, mitochondrial; GROEL; P60 lymphocyte protein; 60 kDa chaperonin; HSP-60; Mitochondrial matrix protein P1; HLD4; SPG13; HSP60; Chaperonin 60; Hsp60; CPN60; HSP65

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:10 - 1:50
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A431	

Properties


Form	Liquid
Purification	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Buffer	PBS and 0.09% (W/V) Sodium azide.
Preservative	0.09% (W/V) Sodium azide.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 3329 Human Swiss-port # P10809 Human
Gene Symbol	HSPD1
Gene Full Name	heat shock 60kDa protein 1 (chaperonin)
Background	This gene encodes a member of the chaperonin family. The encoded mitochondrial protein may function as a signaling molecule in the innate immune system. This protein is essential for the folding and assembly of newly imported proteins in the mitochondria. This gene is adjacent to a related family member and the region between the 2 genes functions as a bidirectional promoter. Several pseudogenes have been associated with this gene. Two transcript variants encoding the same protein have been identified for this gene. Mutations associated with this gene cause autosomal recessive spastic paraplegia 13. [provided by RefSeq, Jun 2010]
Function	Implicated in mitochondrial protein import and macromolecular assembly. May facilitate the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix. [UniProt]
Research Area	Controls and Markers antibody; Signaling Transduction antibody; Mitochondrial Marker antibody
Calculated Mw	61 kDa
Cellular Localization	Mitochondrion matrix.

Images



ARG55686 anti-Hsp 60 antibody ICC/IF image

Immunofluorescence: HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then stained with ARG55686 anti-Hsp 60 antibody (green) at 1:25 dilution, 1 hour at 37°C. Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml, 1 hour at 37°C). DAPI (blue) for nuclear staining.



A431

ARG55686 anti-Hsp 60 antibody WB image

Western blot: 35 µg of A431 cell lysate stained with ARG55686 anti-Hsp 60 antibody.