

Product datasheet

info@arigobio.com

ARG55904 anti-S100A4 antibody

Package: 100 μl, 50 μl Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody recognizes \$100A4

Tested Reactivity Hu

Tested Application IP, WB

Host Mouse

Clonality Monoclonal

Isotype IgG1

Target Name S100A4
Species Human

Immunogen Recombinant Human S100A4 protein.

Conjugation Un-conjugated

Alternate Names S100 calcium-binding protein A4; Placental calcium-binding protein; 42A; MTS1; FSP1; CAPL;

Metastasin; Protein S100-A4; P9KA; 18A2; Protein Mts1; Calvasculin; PEL98

Application Instructions

Application table	Application	Dilution
	IP	Assay-dependent
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Affinity purification with immunogen.

Buffer PBS (pH 7.4), 0.03% Proclin-300 and 50% Glycerol

Preservative 0.03% Proclin-300

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 6275 Human

Swiss-port # P26447 Human

Gene Symbol \$100A4

Gene Full Name S100 calcium binding protein A4

Background The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand

calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in motility, invasion, and tubulin polymerization. Chromosomal rearrangements and altered expression of this gene have been implicated in tumor metastasis. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Jul

2008]

Highlight Related products:


S100A4 antibodies; S100A4 Duos / Panels; Anti-Mouse IgG secondary antibodies;

Related news:

New antibody panels for Myofibroblasts and CAFs

Calculated Mw 12 kDa

Images



ARG55904 anti-S100A4 antibody WB image

Western blot: 30 μg of A549 cell lysate stained with ARG55904 anti-S100A4 antibody at 1:500 dilution.



ARG55904 anti-S100A4 antibody IP image

Immunoprecipitation: HeLa cell lysates were immunoprecipitated and stained with ARG55904 anti-S100A4 antibody.