

## ARG55919 anti-RAGE antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes RAGE
Tested Reactivity	Hu, Ms
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	RAGE
Species	Human
Immunogen	KLH-conjugated synthetic peptide corresponding to aa. 29-59 (N-terminus) of Human RAGE.
Conjugation	Un-conjugated
Alternate Names	Receptor for advanced glycosylation end products; Advanced glycosylation end product-specific receptor; RAGE

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:50 - 1:100
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse lung	

### Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% (W/V) Sodium azide.
Preservative	0.09% (W/V) Sodium azide.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

---

Database links	<a href="#">GeneID: 177 Human</a> <a href="#">Swiss-port # Q15109 Human</a>
Gene Symbol	AGER
Gene Full Name	advanced glycosylation end product-specific receptor
Background	RAGE: Advanced glycosylation end product (AGE) receptor is a member of the immunoglobulin superfamily of cell surface receptors. It is a multiligand receptor, and besides AGE, interacts with other molecules implicated in homeostasis, development, and inflammation, and certain diseases, such as diabetes and Alzheimer's disease. Many alternatively spliced transcript variants encoding different isoforms, as well as non-protein-coding variants, have been described for this gene (PMID:18089847). [provided by RefSeq, May 2011]
Function	RAGE Mediates interactions of advanced glycosylation end products (AGE). These are nonenzymatically glycosylated proteins which accumulate in vascular tissue in aging and at an accelerated rate in diabetes. Acts as a mediator of both acute and chronic vascular inflammation in conditions such as atherosclerosis and in particular as a complication of diabetes. AGE/RAGE signaling plays an important role in regulating the production/expression of TNF-alpha, oxidative stress, and endothelial dysfunction in type 2 diabetes. Interaction with S100A12 on endothelium, mononuclear phagocytes, and lymphocytes triggers cellular activation, with generation of key proinflammatory mediators. Interaction with S100B after myocardial infarction may play a role in myocyte apoptosis by activating ERK1/2 and p53/TP53 signaling. Receptor for amyloid beta peptide. Contributes to the translocation of amyloid-beta peptide (ABPP) across the cell membrane from the extracellular to the intracellular space in cortical neurons. ABPP-initiated RAGE signaling, especially stimulation of p38 mitogen-activated protein kinase (MAPK), has the capacity to drive a transport system delivering ABPP as a complex with RAGE to the intraneuronal space. Can also bind oligonucleotides. [UniProt]
Highlight	Related products: <a href="#">RAGE antibodies</a> ; <a href="#">RAGE ELISA Kits</a> ; <a href="#">Anti-Rabbit IgG secondary antibodies</a> ;
Calculated Mw	43 kDa
Cellular Localization	Isoform 1: Cell membrane; Single-pass type I membrane protein Isoform 10: Cell membrane; Single-pass type I membrane protein


## Images

---



ARG55919 anti-RAGE antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human cancer tissue stained with ARG55919 anti-RAGE antibody.



ARG55919 anti-RAGE antibody WB image

Western blot: 35  $\mu$ g of Mouse lung lysate stained with ARG55919 anti-RAGE antibody.