

## ARG56304 anti-HIF1AN antibody

Package: 50 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes HIF1AN
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	HIF1AN
Species	Human
Immunogen	Recombinant protein of Human HIF1AN
Conjugation	Un-conjugated
Alternate Names	Factor inhibiting HIF-1; FIH-1; Hypoxia-inducible factor asparagine hydroxylase; Hypoxia-inducible factor 1-alpha inhibitor; EC 1.14.11.n4; EC 1.14.11.30; FIH1

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	IP	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	MCF7	


### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation


Database links	<a href="#">GeneID: 319594 Mouse</a> <a href="#">GeneID: 55662 Human</a> <a href="#">Swiss-port # Q8BLR9 Mouse</a> <a href="#">Swiss-port # Q9NWT6 Human</a>
Gene Symbol	HIF1AN
Gene Full Name	hypoxia inducible factor 1, alpha subunit inhibitor
Function	Hydroxylates HIF-1 alpha at 'Asp-803' in the C-terminal transactivation domain (CAD). Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD) of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively regulates NOTCH1 activity, accelerating myogenic differentiation. Positively regulates ASB4 activity, promoting vascular differentiation. [UniProt]
Calculated Mw	40 kDa

## Images



ARG56304 anti-HIF1AN antibody ICC/IF image

Immunofluorescence: U2OS cells stained with ARG56304 anti-HIF1AN antibody.



MCF7

ARG56304 anti-HIF1AN antibody WB image

Western blot: MCF7 cell lysate stained with ARG56304 anti-HIF1AN antibody.