

ARG56516 anti-CD19 antibody [CB19]

Package: 50 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [CB19] recognizes CD19
Tested Reactivity	Hu
Tested Application	FACS, ICC/IF, IHC-Fr
Host	Mouse
Clonality	Monoclonal
Clone	CB19
Isotype	IgG1
Target Name	CD19
Antigen Species	Human
Immunogen	Human chronic lymphocytic leukemia cell line.
Conjugation	Un-conjugated
Alternate Names	Differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte surface antigen B4; B4; CD antigen CD19; CVID3

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	ICC/IF	2 - 10 µg/ml
	IHC-Fr	2 - 10 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Calculated Mw	61 kDa	


Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS (pH 7.2) and 0.02% Sodium azide.
Preservative	0.02% Sodium azide.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 930 Human Swiss-port # P15391 Human
Gene Symbol	CD19
Gene Full Name	CD19 molecule
Background	CD19: Lymphocytes proliferate and differentiate in response to various concentrations of different antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. This gene encodes a cell surface molecule which assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation. [provided by RefSeq, Jul 2008]
Function	CD19 functions as coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes. Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed:2463100, PubMed:1373518, PubMed:16672701). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed:9382888, PubMed:9317126, PubMed:12387743, PubMed:16672701). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:9317126). Required for normal differentiation of B-1 cells. Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:2463100, PubMed:1373518). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed:9317126, PubMed:12387743, PubMed:16672701). [UniProt]
Highlight	Related products: CD19 antibodies ; CD19 Duos / Panels ; Anti-Mouse IgG secondary antibodies ; Related news: Lymphoma

Images



ARG56516 anti-CD19 antibody [CB19] IHC-Fr image

Immunohistochemistry: Frozen sections of small lymphocytic lymphoma (SLL) tissue stained with ARG56516 anti-CD19 antibody [CB19].