

ARG56588 anti-IL4 antibody [1.3_4F11-2G3]

Package: 100 µg, 50 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [1.3_4F11-2G3] recognizes IL4
Tested Reactivity	Hu, Ms
Tested Application	ELISA, WB
Host	Mouse
Clonality	Monoclonal
Clone	1.3_4F11-2G3
Isotype	IgG1, kappa
Target Name	IL4
Species	Human
Immunogen	E.coli derived Recombinant Human IL-4. (MHKCDITLQE IIKTLNSLTE QKTLCTELTV TDIFAASKNT TEKETFCAA TVLRQFYSHH EKDTRCLGAT AQQFHRHKQL IRFLKRLDRN LWGLAGLNSC PVKEANQSTL ENFLERLKI MREKYSKCSS)
Conjugation	Un-conjugated
Alternate Names	BSF-1; IL-4; B-cell stimulatory factor 1; BCGF1; Lymphocyte stimulatory factor 1; BCGF-1; Interleukin-4; Pitracinra; Binetrakin; BSF1

Application Instructions

Application table	Application	Dilution
	ELISA	Sandwich: 0.5 - 1.0 µg/ml with ARG56724 as a detection antibody
	WB	0.20 - 0.50 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 16189 Mouse](#)

[GeneID: 3565 Human](#)

[Swiss-port # P05112 Human](#)

[Swiss-port # P07750 Mouse](#)

Gene Symbol

IL4

Gene Full Name

interleukin 4

Background

The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Function

Participates in at least several B-cell activation processes as well as of other cell types. It is a costimulator of DNA-synthesis. It induces the expression of class II MHC molecules on resting B-cells. It enhances both secretion and cell surface expression of IgE and IgG1. It also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. [UniProt]

Calculated Mw

17 kDa

Images



ARG56588 anti-IL4 antibody [1.3_4F11-2G3] WB image

Western blot: 250 - 0.24 ng (left to right) of Human IL-4 stained with ARG56588 anti-IL4 antibody [1.3_4F11-2G3], under reducing conditions.


Lanes
1 2 3 4 5 6 7 8 9 10 11 12



ARG56588 anti-IL4 antibody [1.3_4F11-2G3] WB image

Western blot: 250 - 0.24 ng (left to right) of Human IL-4 stained with ARG56588 anti-IL4 antibody [1.3_4F11-2G3], under non-reducing conditions.

Lanes
1 2 3 4 5 6 7 8 9 10 11 12



ARG56588 anti-IL4 antibody [1.3_4F11-2G3] standard curve image

Sandwich ELISA: ARG56588 anti-IL4 antibody [1.3_4F11-2G3] as a capture antibody at 0.5 - 1.0 $\mu\text{g}/\text{ml}$ combined with ARG56724 anti-IL4 antibody (Biotin) as a detection antibody at $\sim 2.0 - 4.0 \mu\text{g}/\text{ml}$. Results of a typical standard run with optical density.