

## ARG56648 anti-CCL22 / MDC antibody

Package: 50 µg  
Store at: -20°C

### Summary

|                     |   |
|---------------------|---|
| Product Description | Rabbit Polyclonal antibody recognizes CCL22 / MDC   |
| Tested Reactivity   | Ms  |
| Tested Application  | ELISA, Neut, WB   |
| Host                | Rabbit  |
| Clonality           | Polyclonal  |
| Isotype             | IgG   |
| Target Name         | CCL22 / MDC   |
| Species             | Mouse   |
| Immunogen           | E.coli derived Recombinant Mouse MDC (CCL22).<br>(GPYGANVEDS ICCQDYIRHP LPSRLVKEFF WTSKSCRKPG VVLITVKNRD ICADPRQVWV KLLHKLKLS)  |
| Conjugation         | Un-conjugated   |
| Alternate Names     | CC chemokine STCP-1; Stimulated T-cell chemotactic protein 1; 3-69; Macrophage-derived chemokine; MDC; DC/B-CK; SCYA22; Small-inducible cytokine A22; ABCD-1; 7-69; 1-69; A-152E5.1; STCP-1; 5-69; C-C motif chemokine 22 |

### Application Instructions

| Application table | Application  | Dilution  |
|-------------------|--|---|
|                   | ELISA  | Sandwich: 0.5 - 2.0 µg/ml with ARG56758 as a detection antibody               |
|                   | Neut   | 6 - 9 µg/ml (To yield [ND50] of the biological activity of mMDC (100 ng/ml) ) |
|                   | WB   | 0.1 - 0.2 µg/ml   |
| Application Note  | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. |   |

### Properties


|                     |  |
|---------------------|--|
| Form                | Liquid   |
| Purification        | Affinity purification with immunogen.  |
| Buffer              | PBS (pH 7.2)   |
| Concentration       | 1 mg/ml  |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note                | For laboratory research only, not for drug, diagnostic or other use.   |

## Bioinformation

|                |  |
|----------------|--|
| Database links | <a href="#">GeneID: 20299 Mouse</a><br><a href="#">Swiss-port # O88430 Mouse</a>   |
| Gene Symbol    | Ccl22  |
| Gene Full Name | chemokine (C-C motif) ligand 22  |
| Background     | This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for primary activated T lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T lymphocytes. The product of this gene binds to chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated T lymphocytes to inflammatory sites and other aspects of activated T lymphocyte physiology. [provided by RefSeq, Sep 2014] |
| Function       | May play a role in the trafficking of activated/effector T-lymphocytes to inflammatory sites and other aspects of activated T-lymphocyte physiology. Chemotactic for monocytes, dendritic cells and natural killer cells. Mild chemoattractant for primary activated T-lymphocytes and a potent chemoattractant for chronically activated T-lymphocytes but has no chemoattractant activity for neutrophils, eosinophils, and resting T-lymphocytes. Binds to CCR4. Processed forms MDC(3-69), MDC(5-69) and MDC(7-69) seem not be active. [UniProt]   |
| Calculated Mw  | 11 kDa   |
| PTM            | The N-terminal processed forms MDC(3-69), MDC(5-69) and MDC(7-69) are produced by proteolytic cleavage after secretion from monocyte derived dendrocytes.  |


## Images

ARG56648 anti-CCL22 / MDC antibody WB image




Western blot: 250 - 0.24 ng of Mouse MDC stained with ARG56648 anti-CCL22 / MDC antibody, under reducing conditions.

ARG56648 anti-CCL22 / MDC antibody WB image



Western blot: 250 - 0.24 ng of Mouse MDC stained with ARG56648 anti-CCL22 / MDC antibody, under non-reducing conditions.



### ARG56648 anti-CCL22 / MDC antibody standard curve image

Sandwich ELISA: ARG56648 anti-CCL22 / MDC antibody as a capture antibody at 0.5 - 2.0 µg/ml combined with ARG56758 anti-CCL22 / MDC antibody (Biotin) as a detection antibody. Results of a typical standard run with optical density.