

ARG56771 anti-CXCL2 / MIP2 antibody (Biotin)

Package: 50 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes CXCL2 / MIP2
Tested Reactivity	Ms
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CXCL2 / MIP2
Species	Mouse
Immunogen	E.coli derived Recombinant Mouse MIP-2 (CXCL2). (AVVASELRQC CLKTLPRVDF KNIQSLSVTP GPGHCAQTEV IATLKGGQKV CLDPEAPLVQ KIIQKILNKG KAN)
Conjugation	Biotin
Alternate Names	Gro-beta; SCYB2; HSF; CINC-2a; GROb; MGSA-b; SB-251353; MIP2A; MIP2; Hematopoietic synergistic factor; 5-73; C-X-C motif chemokine 2; MIP-2a; GRO2; Macrophage inflammatory protein 2-alpha; GRO-beta-T; Growth-regulated protein beta; MIP2-alpha

Application Instructions


Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG56661 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Database links	GeneID: 20310 Mouse Swiss-port # P10889 Mouse
Gene Symbol	Cxcl2
Gene Full Name	chemokine (C-X-C motif) ligand 2
Background	This antimicrobial gene is part of a chemokine superfamily that encodes secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CXC subfamily, is expressed at sites of inflammation and may suppress hematopoietic progenitor cell proliferation. [provided by RefSeq, Sep 2014]
Function	Produced by activated monocytes and neutrophils and expressed at sites of inflammation. Hematoregulatory chemokine, which, in vitro, suppresses hematopoietic progenitor cell proliferation. GRO-beta(5-73) shows a highly enhanced hematopoietic activity. [UniProt]
Calculated Mw	11 kDa
PTM	The N-terminal processed form GRO-beta(5-73) is produced by proteolytic cleavage after secretion from bone marrow stromal cells.

Images



ARG56771 anti-CXCL2 / MIP2 antibody (Biotin) standard curve image

Direct ELISA: ARG56771 anti-CXCL2 / MIP2 antibody (Biotin) at 0.25 - 1.0 µg/ml results of a typical standard run with optical density.



ARG56771 anti-CXCL2 / MIP2 antibody (Biotin) standard curve image

Sandwich ELISA: ARG56771 anti-CXCL2 / MIP2 antibody (Biotin) as a detection antibody at 0.25 - 1.0 µg/ml combined with ARG56661 anti-CXCL2 / MIP-2 antibody as a capture antibody. Results of a typical standard run with optical density.