

## ARG56793 anti-IL7 antibody (Biotin)

Package: 50 µg  
Store at: 4°C

### Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes IL7
Tested Reactivity	Rat
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	IL7
Species	Rat
Immunogen	E.coli derived Recombinant Rat IL-7. (MDCHIKDKDG KAFGSVLMIS INQLDKMTGT DSDCPNNEPN FFKKHLCDT KEAAFLNRRAA RKLRLQFLKMN ISEEFNDHLL RVSDGTQTLV NCTSKEEKT KEQKKNDPCF LKRLLEIKT CWNKILKGS)
Conjugation	Biotin
Alternate Names	IL-7; Interleukin-7

### Application Instructions

Application table	Application	Dilution
	ELISA	Direct: ~ 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG56684 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Database links	<a href="#">GeneID: 25647 Rat</a>
Gene Symbol	IL7
Gene Full Name	interleukin 7
Background	The protein encoded by this gene is a cytokine important for B and T cell development. This cytokine and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. This cytokine is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRB) during early T cell development. This cytokine can be produced locally by intestinal epithelial and epithelial goblet cells, and may serve as a regulatory factor for intestinal mucosal lymphocytes. Knockout studies in mice suggested that this cytokine plays an essential role in lymphoid cell survival. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional splice variants have been described but their presence in normal tissues has not been confirmed.[provided by RefSeq, Dec 2010]
Function	Hematopoietic growth factor capable of stimulating the proliferation of lymphoid progenitors. It is important for proliferation during certain stages of B-cell maturation. [UniProt]
Calculated Mw	20 kDa

## Images



ARG56793 anti-IL7 antibody (Biotin) WB image

Western blot: 250 - 0.24 ng of Rat IL-7 stained with ARG56793 anti-IL7 antibody (Biotin), under reducing conditions.



ARG56793 anti-IL7 antibody (Biotin) WB image

Western blot: 250 - 0.24 ng of Rat IL-7 stained with ARG56793 anti-IL7 antibody (Biotin), under non-reducing conditions.