

ARG56806 anti-IL17 / IL17A antibody (Biotin)

Package: 50 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Goat Polyclonal antibody recognizes IL17 / IL17A
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IL17 / IL17A
Species	Human
Immunogen	E.coli derived Recombinant Human IL-17 (IL-17A). (MIVKAGITIP RNP GCPNSED KNFPRTVMVN LNIHNRNTNT NPKRSSDYNN RSTSPWNLHR NEDPERYPSV IWEAKCRHLG CINADGNVDY HMNSVPIQQE ILVLRREPPH CPNSFRLEKI LVSVGCTCVT PIVHHVA)
Conjugation	Biotin
Alternate Names	IL-17A; Interleukin-17A; IL-17; CTLA8; IL17; CTLA-8; Cytotoxic T-lymphocyte-associated antigen 8

Application Instructions

Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml as a detection antibody and with ARG56697 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	Sandwich ELISA: Sandwich ELISA: Recommended antibody pairs in sandwich immunoassay for IL17 / IL17A (capture - detection): ARG56697 - ARG56806 * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 3605 Human Swiss-port # Q16552 Human
Gene Symbol	IL17A
Gene Full Name	interleukin 17A
Background	The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. [provided by RefSeq, Jul 2008]
Function	Induces stromal cells to produce proinflammatory and hematopoietic cytokines. Enhances the surface expression of ICAM1/intracellular adhesion molecule 1 in fibroblasts. [UniProt]
Highlight	Related products: IL17 antibodies ; IL17 ELISA Kits ; IL17 recombinant proteins ; Anti-Goat IgG secondary antibodies ; Related news: HMGB1, a biomarker and therapeutic target in COVID-19
Calculated Mw	18 kDa
PTM	Found both in glycosylated and nonglycosylated forms.

Images



ARG56806 anti-IL17 / IL17A antibody (Biotin) standard curve image

Direct ELISA: ARG56806 anti-IL17 / IL17A antibody (Biotin) at 0.25 - 1.0 µg/ml results of a typical standard run with optical density.



ARG56806 anti-IL17 / IL17A antibody (Biotin) standard curve image

Sandwich ELISA: ARG56806 anti-IL17 / IL17A antibody (Biotin) as a detection antibody at 0.25 - 1.0 µg/ml combined with ARG56697 anti-IL17 / IL17A antibody as a capture antibody. Results of a typical standard run with optical density.