

## ARG57599 anti-STAT2 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes STAT2
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	STAT2
Species	Human
Immunogen	Synthetic peptide derived from Human Stat2.
Conjugation	Un-conjugated
Alternate Names	P113; Signal transducer and activator of transcription 2; STAT113; p113; ISGF-3

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:1000 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	STAT2
Gene Full Name	signal transducer and activator of transcription 2, 113kDa
Background	<p>The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. In response to interferon (IFN), this protein forms a complex with STAT1 and IFN regulatory factor family protein p48 (ISGF3G), in which this protein acts as a transactivator, but lacks the ability to bind DNA directly. Transcription adaptor P300/CBP (EP300/CREBBP) has been shown to interact specifically with this protein, which is thought to be involved in the process of blocking IFN-alpha response by adenovirus. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2010]</p>
Function	<p>Signal transducer and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with IRF9/ISGF3G to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state. [UniProt]</p>
Highlight	<p>Related products: <a href="#">STAT2 antibodies</a>; <a href="#">Anti-Rabbit IgG secondary antibodies</a>; Related news: <a href="#">Exploring Antiviral Immune Response</a> <a href="#">circNDUFB2, a circular RNA (circRNA), activates anti-tumor immunity</a></p>
Calculated Mw	98 kDa
PTM	<p>Tyrosine phosphorylated in response to IFN-alpha. Phosphorylation at Ser-287 negatively regulates the transcriptional response. [UniProt]</p>
Cellular Localization	Late endosome membrane > Peripheral membrane protein

## Images




ARG57599 anti-STAT2 antibody ICC/IF image

Immunofluorescence: A431 cells stained with ARG57599 anti-STAT2 antibody.



ARG57599 anti-STAT2 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human lung cancer tissue stained with ARG57599 anti-STAT2 antibody.



K562

ARG57599 anti-STAT2 antibody WB image

Western blot: K562 cell lysate stained with ARG57599 anti-STAT2 antibody.