

## ARG57827 anti-14-3-3 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes 14-3-3
Tested Reactivity	Hu
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	14-3-3
Species	Human
Immunogen	KLH-conjugated synthetic peptide within aa. 217-246 of Human 14-3-3.
Conjugation	Un-conjugated
Alternate Names	Protein 1054; 14-3-3 protein beta/alpha; Protein kinase C inhibitor protein 1; HS1; GW128; KCIP-1; YWHAA; HEL-S-1

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:10 - 1:50
	IHC-P	1:50 - 1:100
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Jurkat	
Observed Size	~ 28 kDa	


### Properties

Form	Liquid
Purification	Saturated Ammonium Sulfate precipitation followed by dialysis against PBS.
Buffer	PBS and 0.09% (W/V) Sodium azide.
Preservative	0.09% (W/V) Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation


Gene Symbol	YWHAB
Gene Full Name	tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, beta
Background	This gene encodes a protein belonging to the 14-3-3 family of proteins, members of which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals. The encoded protein has been shown to interact with RAF1 and CDC25 phosphatases, suggesting that it may play a role in linking mitogenic signaling and the cell cycle machinery. Two transcript variants, which encode the same protein, have been identified for this gene. [provided by RefSeq, Jul 2008]
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner. Negative regulator of osteogenesis. Blocks the nuclear translocation of the phosphorylated form (by AKT1) of SRPK2 and antagonizes its stimulatory effect on cyclin D1 expression resulting in blockage of neuronal apoptosis elicited by SRPK2. [UniProt]
Calculated Mw	28 kDa
PTM	The alpha, brain-specific form differs from the beta form in being phosphorylated. Phosphorylated on Ser-60 by protein kinase C delta type catalytic subunit in a sphingosine-dependent fashion. [UniProt]
Cellular Localization	Cytoplasm. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. [UniProt]

## Images



ARG57827 anti-14-3-3 antibody ICC/IF image


Immunofluorescence: HeLa cells stained with ARG57827 anti-14-3-3 antibody (green). DAPI (blue) for nuclear staining.



ARG57827 anti-14-3-3 antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human breast carcinoma tissue stained with ARG57827 anti-14-3-3 antibody.

ARG57827 anti-14-3-3 antibody WB image



Western blot: Jurkat cell lysate stained with ARG57827 anti-14-3-3 antibody.

Jurkat

---