

ARG58064 anti-Mesothelin antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Mesothelin
Tested Reactivity	Hu
Tested Application	IHC-P, IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Mesothelin
Species	Human
Immunogen	Synthetic peptide derived from Human Mesothelin.
Conjugation	Un-conjugated
Alternate Names	MPF; Mesothelin; CAK1 antigen; Pre-pro-megakaryocyte-potentiating factor; SMRP

Application Instructions

Application table	Application	Dilution
	IHC-P	1:100 - 1:500
	IP	1:50
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	
Observed Size	~ 69 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation


Gene Symbol	MSLN
Gene Full Name	mesothelin
Background	This gene encodes a precursor protein that is cleaved into two products, megakaryocyte potentiating factor and mesothelin. Megakaryocyte potentiation factor functions as a cytokine that can stimulate colony formation in bone marrow megakaryocytes. Mesothelian is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed in epithelial mesotheliomas, ovarian cancers and in specific squamous cell carcinomas. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Apr 2010]
Function	Membrane-anchored forms may play a role in cellular adhesion. Megakaryocyte-potentiating factor (MPF) potentiates megakaryocyte colony formation in vitro. [UniProt]
Calculated Mw	69 kDa
PTM	Both MPF and the cleaved form of mesothelin are N-glycosylated. Proteolytically cleaved by a furin-like convertase to generate megakaryocyte-potentiating factor (MPF), and the cleaved form of mesothelin. [UniProt]

Images



ARG58064 anti-Mesothelin antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human tonsil stained with ARG58064 anti-Mesothelin antibody at 1:500 dilution.



ARG58064 anti-Mesothelin antibody WB image

Western blot: HeLa cell lysate stained with ARG58064 anti-Mesothelin antibody.