

## ARG58145 anti-delta 1 Catenin antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes delta 1 Catenin
Tested Reactivity	Hu, Ms
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	delta 1 Catenin
Species	Human
Immunogen	Synthetic peptide derived from Human delta 1 Catenin.
Conjugation	Un-conjugated
Alternate Names	p120 catenin; p120; CAS; P120CAS; CTNND; p120(CAS); cas; Catenin delta-1; Cadherin-associated Src substrate; p120(CTN); P120CTN; ctn

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:100
	IHC-P	1:50 - 1:100
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	


### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	CTNND1
Gene Full Name	catenin (cadherin-associated protein), delta 1
Background	This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene. [provided by RefSeq, Dec 2010]
Function	Binds to and inhibits the transcriptional repressor ZBTB33, which may lead to activation of target genes of the Wnt signaling pathway (By similarity). Associates with and regulates the cell adhesion properties of both C-, E- and N-cadherins, being critical for their surface stability. Implicated both in cell transformation by SRC and in ligand-induced receptor signaling through the EGF, PDGF, CSF-1 and ERBB2 receptors. Promotes GLIS2 C-terminal cleavage. [UniProt]
Calculated Mw	108 kDa
PTM	Phosphorylated by FER and other protein-tyrosine kinases. Phosphorylated at Ser-288 by PAK5. Dephosphorylated by PTPRJ. [UniProt]
Cellular Localization	Nucleus. [UniProt]

## Images



ARG58145 anti-delta 1 Catenin antibody ICC/IF image

Immunofluorescence: MCF-7 cells stained with ARG58145 anti-delta 1 Catenin antibody.



ARG58145 anti-delta 1 Catenin antibody WB image

Western blot: HeLa cell lysate stained with ARG58145 anti-delta 1 Catenin antibody.