

Product datasheet

info@arigobio.com

ARG58421 anti-CHRNA10 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes CHRNA10

Tested Reactivity Hu
Predict Reactivity Rat

Tested Application IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name CHRNA10
Species Human

Immunogen KLH-conjugated synthetic peptide corresponding to aa. 179-206 (Center) of Human CHRNA10.

Conjugation Un-conjugated

Alternate Names NACHR alpha-10; Neuronal acetylcholine receptor subunit alpha-10; Nicotinic acetylcholine receptor

subunit alpha-10

Application Instructions

Application table	Application	Dilution
	IHC-P	1:10 - 1:50
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	NCI-H292	

Properties

Form Liquid

Purification Purification with Protein A and immunogen peptide.

Buffer PBS and 0.09% (W/V) Sodium azide.

Preservative 0.09% (W/V) Sodium azide.

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CHRNA10

Gene Full Name cholinergic receptor, nicotinic, alpha 10 (neuronal)

Function Ionotropic receptor with a probable role in the modulation of auditory stimuli. Agonist binding may

induce an extensive change in conformation that affects all subunits and leads to opening of an ion-conducting channel across the plasma membrane. The channel is permeable to a range of divalent cations including calcium, the influx of which may activate a potassium current which hyperpolarizes the cell membrane. In the ear, this may lead to a reduction in basilar membrane motion, altering the activity of auditory nerve fibers and reducing the range of dynamic hearing. This may protect against


acoustic trauma. [UniProt]

Calculated Mw 50 kDa

Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein Cell membrane;

Multi-pass membrane protein. [UniProt]

Images



ARG58421 anti-CHRNA10 antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human tonsil tissue stained with ARG58421 anti-CHRNA10 antibody.



ARG58421 anti-CHRNA10 antibody WB image

Western blot: 35 μg of NCI-H292 cell lysate stained with ARG58421 anti-CHRNA10 antibody.

NCI-H292