

Product datasheet

info@arigobio.com

ARG58598 anti-FGF8 antibody

Package: 100 μl Store at: -20°C

Summary

Host

Product Description Rabbit Polyclonal antibody recognizes FGF8

Rabbit

Tested Reactivity Hu
Tested Application WB

Clonality Polyclonal

Isotype IgG

Target Name FGF8

Species Human

Immunogen KLH-conjugated synthetic peptide between aa. 32-66 of Human FGF8.

Conjugation Un-conjugated

Alternate Names Fibroblast growth factor 8; HH6; AIGF; FGF-8; HBGF-8; KAL6; Heparin-binding growth factor 8;

Androgen-induced growth factor

Application Instructions

Application table	Application	Dilution
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	LNCaP	

Properties

Form Liquid

Purification Purification with Protein A and immunogen peptide.

Buffer PBS and 0.09% (W/V) sodium azide.

Preservative 0.09% (W/V) Sodium azide

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol FGF8

Gene Full Name fibroblast growth factor 8 (androgen-induced)

Background The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family

members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is known to be a factor that supports androgen and anchorage independent growth of mammary tumor cells. Overexpression of this gene has been shown to increase tumor growth and angiogensis. The adult expression of this gene is restricted to testes and ovaries. Temporal and spatial pattern of this gene expression suggests its function as an embryonic epithelial factor. Studies of the mouse and chick homologs revealed roles in midbrain and limb development, organogenesis, embryo gastrulation and left-right axis determination. The alternative splicing of this

gene results in four transcript variants. [provided by RefSeq, Jul 2008]

Function Plays an important role in the regulation of embryonic development, cell proliferation, cell

differentiation and cell migration. Required for normal brain, eye, ear and limb development during embryogenesis. Required for normal development of the gonadotropin-releasing hormone (GnRH)

neuronal system. Plays a role in neurite outgrowth in hippocampal cells. [UniProt]

Calculated Mw 27 kDa

Cellular Localization Secreted. [UniProt]

Images

