

ARG58704 anti-FMO5 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes FMO5
Tested Reactivity	Hu, Ms
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	FMO5
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 77-90 of Human FMO5 (DHYPNFMHNAQVLE).
Conjugation	Un-conjugated
Alternate Names	Dimethylaniline monooxygenase [N-oxide-forming] 5; FMO 5; Hepatic flavin-containing monooxygenase 5; EC 1.14.13.8; Dimethylaniline oxidase 5

Application Instructions

Application table	<table> <tr> <th>Application</th><th>Dilution</th></tr> <tr> <td>WB</td><td>0.1 - 0.5 µg/ml</td></tr> </table>	Application	Dilution	WB	0.1 - 0.5 µg/ml
Application	Dilution				
WB	0.1 - 0.5 µg/ml				
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.9% NaCl, 0.2% Na2HPO4, 0.05% Thimerosal, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Thimerosal and 0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	FMO5
Gene Full Name	flavin containing monooxygenase 5
Background	Metabolic N-oxidation of the diet-derived amino-trimethylamine (TMA) is mediated by flavin-containing monooxygenase and is subject to an inherited FMO3 polymorphism in man resulting in a small subpopulation with reduced TMA N-oxidation capacity resulting in fish odor syndrome Trimethylaminuria. Three forms of the enzyme, FMO1 found in fetal liver, FMO2 found in adult liver, and FMO3 are encoded by genes clustered in the 1q23-q25 region. Flavin-containing monooxygenases are NADPH-dependent flavoenzymes that catalyzes the oxidation of soft nucleophilic heteroatom centers in drugs, pesticides, and xenobiotics. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2009]
Function	In contrast with other forms of FMO it does not seem to be a drug-metabolizing enzyme. [UniProt]
Calculated Mw	60 kDa
Cellular Localization	Microsome membrane. Endoplasmic reticulum membrane. [UniProt]

Images



ARG58704 anti-FMO5 antibody WB image

Western blot: 50 µg of Mouse liver, Mouse testis and Mouse spleen lysates stained with ARG58704 anti-FMO5 antibody at 0.5 µg/ml dilution.



ARG58704 anti-FMO5 antibody WB image

Western blot: 50 µg of sample under reducing conditions. HeLa, Mouse liver, Mouse testis, Mouse spleen lysates stained with ARG58704 anti-FMO5 antibody at 0.5 µg/ml dilution, overnight at 4°C.