

# Product datasheet

info@arigobio.com

# ARG58996 anti-GDI2 antibody

Package: 100 μl Store at: -20°C

#### **Summary**

Product Description Rabbit Polyclonal antibody recognizes GDI2

Tested Reactivity Hu

Tested Application IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name GDI2

Species Human

Immunogen KLH-conjugated synthetic peptide corresponding to aa. 135-164 of Human GDI2.

Conjugation Un-conjugated

Alternate Names HEL-S-46e; Rab GDP dissociation inhibitor beta; Rab GDI beta; RABGDIB; Guanosine diphosphate

dissociation inhibitor 2; GDI-2

## **Application Instructions**

Application table	Application	Dilution
	IHC-P	1:10 - 1:50
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	MCF7	

#### **Properties**

Form Liquid

Purification Purification with Protein A and immunogen peptide.

Buffer PBS and 0.09% (W/V) Sodium azide.

Preservative 0.09% (W/V) Sodium azide

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol GDI2

Gene Full Name GDP dissociation inhibitor 2

Background GDP dissociation inhibitors are proteins that regulate the GDP-GTP exchange reaction of members of

the rab family, small GTP-binding proteins of the ras superfamily, that are involved in vesicular trafficking of molecules between cellular organelles. GDIs slow the rate of dissociation of GDP from rab proteins and release GDP from membrane-bound rabs. GDI2 is ubiquitously expressed. The GDI2 gene

contains many repetitive elements indicating that it may be prone to inversion/deletion

rearrangements. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

[provided by RefSeq, Jul 2008]


Function Regulates the GDP/GTP exchange reaction of most Rab proteins by inhibiting the dissociation of GDP

from them, and the subsequent binding of GTP to them. [UniProt]

Calculated Mw 51 kDa

Cellular Localization Cytoplasm. Membrane; Peripheral membrane protein. [UniProt]

### **Images**



#### ARG58996 anti-GDI2 antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human liver tissue stained with ARG58996 anti-GDI2 antibody.



**-** 35

**-** 25

**-** 16

#### ARG58996 anti-GDI2 antibody WB image

Western blot: 35  $\mu g$  of MCF7 cell lysate stained with ARG58996 anti-GDI2 antibody.

MCF7