

ARG59173 anti-HDAC9 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes HDAC9
Tested Reactivity	Hu, Ms
Predict Reactivity	Chk
Tested Application	ICC/IF, IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	HDAC9
Species	Human
Immunogen	KLH-conjugated synthetic peptide corresponding to aa. 2-32 of Human HDAC9.
Conjugation	Un-conjugated
Alternate Names	HD7; MEF2-interacting transcription repressor MITR; Histone deacetylase-related protein; HDAC9B; Histone deacetylase 9; HDAC7; HDRP; HDAC9FL; MITR; HDAC7B; HD9; HD7b; EC 3.5.1.98; HDAC; Histone deacetylase 7B

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:1000
	IP	1:100
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% (W/V) Sodium azide.
Preservative	0.09% (W/V) Sodium azide.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	HDAC9
Gene Full Name	histone deacetylase 9
Background	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene has sequence homology to members of the histone deacetylase family. This gene is orthologous to the Xenopus and mouse MITR genes. The MITR protein lacks the histone deacetylase catalytic domain. It represses MEF2 activity through recruitment of multicomponent corepressor complexes that include CtBP and HDACs. This encoded protein may play a role in hematopoiesis. Multiple alternatively spliced transcripts have been described for this gene but the full-length nature of some of them has not been determined. [provided by RefSeq, Jul 2008]
Function	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses MEF2-dependent transcription. Isoform 3 lacks active site residues and therefore is catalytically inactive. Represses MEF2-dependent transcription by recruiting HDAC1 and/or HDAC3. Seems to inhibit skeletal myogenesis and to be involved in heart development. Protects neurons from apoptosis, both by inhibiting JUN phosphorylation by MAPK10 and by repressing JUN transcription via HDAC1 recruitment to JUN promoter. [UniProt]
Calculated Mw	111 kDa
PTM	Phosphorylated on Ser-220 and Ser-450; which promotes 14-3-3-binding, impairs interaction with MEF2, and antagonizes antimyogenic activity. Phosphorylated on Ser-240; which impairs nuclear accumulation (By similarity). Isoform 7 is phosphorylated on Tyr-1010. Phosphorylated by the PKC kinases PKN1 and PKN2, impairing nuclear import. Sumoylated. [UniProt]
Cellular Localization	Nucleus. [UniProt]

Images



ARG59173 anti-HDAC9 antibody ICC/IF image

Immunofluorescence: Undifferentiated C2C12 myoblasts transfected with a HDAC9-expressing plasmid. HDAC9 is detected by using ARG59173 anti-HDAC9 antibody.