

## ARG59225 anti-SMC3 antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes SMC3
Tested Reactivity	Hu, Ms, Rat
Predict Reactivity	Bov, Chk, Hrs, Mk, Rb
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	SMC3
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 1178-1216 of Human SMC3. (ELLESADKFGVVKFRNKVSHIDVITAEMAKDFVEDDTTH)
Conjugation	Un-conjugated
Alternate Names	CDLS3; CSPG6; SMC3L1; SMC-3; Chromosome-associated polypeptide; BMH; Basement membrane-associated chondroitin proteoglycan; Structural maintenance of chromosomes protein 3; Bamacan; hCAP; Chondroitin sulfate proteoglycan 6; SMC protein 3; HCAP; BAM

### Application Instructions

Application table	Application	Dilution
	WB	0.1 - 0.5 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Gene Symbol	SMC3
Gene Full Name	structural maintenance of chromosomes 3
Background	This gene belongs to the SMC3 subfamily of SMC proteins. The encoded protein occurs in certain cell types as either an intracellular, nuclear protein or a secreted protein. The nuclear form, known as structural maintenance of chromosomes 3, is a component of the multimeric cohesin complex that holds together sister chromatids during mitosis, enabling proper chromosome segregation. Post-translational modification of the encoded protein by the addition of chondroitin sulfate chains gives rise to the secreted proteoglycan bamacan, an abundant basement membrane protein. [provided by RefSeq, Jul 2008]
Function	Central component of cohesin, a complex required for chromosome cohesion during the cell cycle. The cohesin complex may form a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. Cohesion is coupled to DNA replication and is involved in DNA repair. The cohesin complex plays also an important role in spindle pole assembly during mitosis and in chromosomes movement. [UniProt]
Calculated Mw	142 kDa
PTM	Phosphorylated at Ser-1083 in a SPO11-dependent manner.  Acetylation at Lys-105 and Lys-106 by ESCO1 is important for genome stability and S phase sister chromatid cohesion. Regulated by DSCC1, it is required for processive DNA synthesis, coupling sister chromatid cohesion establishment during S phase to DNA replication. Deacetylation by HDAC8, regulates release of the cohesin complex from chromatin. [UniProt]
Cellular Localization	Nucleus. Chromosome. Chromosome, centromere. Note=Associates with chromatin. [UniProt]

## Images



ARG59225 anti-SMC3 antibody WB image

Western blot: 50 µg of Rat brain, 50 µg of Rat liver, 50 µg of Rat testis, 40 µg of HeLa, 40 µg of A549, 40 µg of MCF-7 and 40 µg of NIH/3T3 lysates stained with ARG59225 anti-SMC3 antibody at 0.5 µg/ml dilution.