

## ARG59965 anti-RAB31 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes RAB31
Tested Reactivity	Hu, Ms, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	RAB31
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 1-195 of Human RAB31 (NP_006859.2).
Conjugation	Un-conjugated
Alternate Names	Rab22B; Ras-related protein Rab-22B; Ras-related protein Rab-31

### Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Rat lung, Mouse lung and HL-60	
Observed Size	22 kDa	


### Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	RAB31
Gene Full Name	RAB31, member RAS oncogene family
Background	Small GTP-binding proteins of the RAB family, such as RAB31, play essential roles in vesicle and granule targeting (Bao et al., 2002 [PubMed 11784320]).[supplied by OMIM, Jul 2009]
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. Required for the integrity and for normal function of the Golgi apparatus and the trans-Golgi network. Plays a role in insulin-stimulated translocation of GLUT4 to the cell membrane. Plays a role in M6PR transport from the trans-Golgi network to endosomes. Plays a role in the internalization of EGFR from the cell membrane into endosomes. Plays a role in the maturation of phagosomes that engulf pathogens, such as S.aureus and M.tuberculosis. [UniProt]
Calculated Mw	22 kDa
Cellular Localization	Golgi apparatus, trans-Golgi network. Golgi apparatus, trans-Golgi network membrane; Lipid-anchor; Cytoplasmic side. Early endosome. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Note=Rapidly recruited to phagosomes containing S.aureus or M.tuberculosis (PubMed:21255211). [UniProt]

Images



ARG59965 anti-RAB31 antibody WB image

Western blot: 25 µg of Rat lung, Mouse lung and HL-60 cell lysates stained with ARG59965 anti-RAB31 antibody at 1:1000 dilution.