

ARG62702 anti-CD105 / Endoglin antibody [MEM-229] (FITC)

Package: 100 tests
Store at: 4°C

Summary

| | |
|---------------------|---|
| Product Description | FITC-conjugated Mouse Monoclonal antibody [MEM-229] recognizes CD105 / Endoglin |
| Tested Reactivity | Hu, Pig |
| Tested Application | FACS |
| Specificity | The clone MEM-229 recognizes CD105 (Endoglin), a 90 kDa type I integral membrane homodimer glycoprotein expressed on vascular endothelial cells (small and large vessels), activated monocytes and tissue macrophages, stromal cells of certain tissues including bone marrow, pre-B lymphocytes in fetal marrow and erythroid precursors in fetal and adult bone marrow; it is also present on syncytiotrophoblast on placenta throughout pregnancy. |
| Host | Mouse |
| Clonality | Monoclonal |
| Clone | MEM-229 |
| Isotype | IgG2a |
| Target Name | CD105 / Endoglin |
| Species | Human |
| Immunogen | Recombinant Vaccinia virus containing the human CD105 (L-isoform) cDNA. |
| Conjugation | FITC |
| Alternate Names | CD antigen CD105; HHT1; Endoglin; ORW1; END |

Application Instructions

| Application table | Application | Dilution |
|-------------------|-------------|-------------------------------|
| | FACS | 20 µl / 10 ⁶ cells |

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification Note | The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary. |
| Buffer | PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA |
| Preservative | 15 mM Sodium azide |
| Stabilizer | 0.2% (w/v) high-grade protease free BSA |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

| | |
|----------------|---|
| Database links | GeneID: 2022 Human GeneID: 397096 Pig Swiss-port # P17813 Human Swiss-port # P37176 Pig |
| Gene Symbol | ENG |
| Gene Full Name | endoglin |
| Background | CD105 (Endoglin) is a homodimeric transmembrane glycoprotein serving in presence of TGFbetaR-2 as a receptor for TGFbeta-1 and TGFbeta-3. CD105 is highly expressed on endothelial cells and promotes angiogenesis during wound healing, infarcts and in a wide range of tumours and its gene expression is stimulated by hypoxia. CD105 prevents apoptosis in hypoxic endothelial cells and also antagonises the inhibitory effects of TGFbeta-1 on vascular endothelial cell growth and migration. Normal cellular levels of CD105 are required for formation of new blood vessels. |
| Function | Major glycoprotein of vascular endothelium. Involved in the regulation of angiogenesis. May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors. Acts as TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade. Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells and modulates TGF-beta1 signaling through SMAD3. [UniProt] |
| Research Area | Cancer antibody; Cell Biology and Cellular Response antibody; Controls and Markers antibody; Developmental Biology antibody; Immune System antibody |
| Calculated Mw | 71 kDa |